

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
16 June 2005 (16.06.2005)

PCT

(10) International Publication Number  
**WO 2005/054199 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 213/74**,  
237/20, 239/42, 241/20, 253/07, 401/12, 401/14, 413/10,  
403/10, A61K 31/4418, 31/444, 31/496, 31/497, 31/506,  
31/53, 31/5377, A61P 29/00, 37/00

Street, ELWOOD, VIC 3184 (AU). SIMS, Colette, Gloria  
[AU/AU]; 191 Kew Street, KEWDALE, W.A. 6105 (AU).

(21) International Application Number:  
PCT/AU2004/001689

(74) Agent: **BLAKE DAWSON WALDRON PATENT SER-  
VICES**; level 39, 101 Collins Street, MELBOURNE, VIC  
3000 (AU).

(22) International Filing Date: 3 December 2004 (03.12.2004)

(81) Designated States (*unless otherwise indicated, for every  
kind of national protection available*): AE, AG, AL, AM,  
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,  
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,  
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,  
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,  
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,  
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,  
ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
2003906680 3 December 2003 (03.12.2003) AU

(84) Designated States (*unless otherwise indicated, for every  
kind of regional protection available*): ARIPO (BW, GH,  
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,  
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,  
FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO,  
SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN,  
GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **CY-  
TOPIA RESEARCH PTY LTD** [AU/AU]; Level 5,  
Baker Heart Research Institute Building, MELBOURNE,  
VIC 3004 (AU).

(72) Inventors; and





(75) Inventors/Applicants (*for US only*): **BURNS, Christo-  
pher, John** [AU/AU]; 3 Browning Street, SEDDON,  
VIC 3011 (AU). **WILKS, Andrew, Frederick** [AU/AU];  
6 MacFarlan Lane, SOUTH YARRA, VIC 3141 (AU).  
**HARTE, Michael, Francis** [AU/AU]; 9/62 Meyrick  
Crescent, VIEWBANK, VIC 3084 (AU). **SIKANYIKA,  
Harrison** [ZM/AU]; 2 Pitt Road, KILSYTH, VIC 3137  
(AU). **FANTINO, Emmanuelle** [FR/AU]; 1/3 Alfriston

Published:

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

(54) Title: TUBULIN INHIBITORS



(57) Abstract: Compounds of general formula (I), (II) (III) and (V) are described for use in modulating microtubule polymerisation and in the treatment of associated disease states. Use of compounds (I), (III) and (V) in the treatment of kinase-associated disease states is also described. Further described are novel compounds of formula (II), (III) and (V).

WO 2005/054199 A1

1.

## TUBULIN INHIBITORS

### FIELD OF THE INVENTION

5 The present invention relates to methods of suppressing the growth of cancers and other proliferative diseases by the administration of compounds that act by binding to tubulin and to novel heterocyclic compounds suitable for such administration.

### BACKGROUND OF THE INVENTION

10 There are many human and veterinary diseases that stem from processes of uncontrolled or abnormal cellular proliferation. Most important among these diseases is cancer, the generic name given to a broad range of cellular malignancies characterized by unregulated growth and lack of differentiation. Psoriasis is another disease that is characterized by uncontrolled or abnormal cellular proliferation. Psoriasis is a common chronic skin disease characterized by the presence of dry scales and plaques. The disease results from hyperproliferation of the epidermis and incomplete differentiation of  
15 keratinocytes. Psoriasis often involves the scalp, elbows, knees, back, buttocks, nails, eyebrows, and genital regions, and may range in severity from mild to extremely debilitating, resulting in psoriatic arthritis, pustular psoriasis, and exfoliative psoriatic dermatitis. There is, at present, no general therapeutic cure that exists for psoriasis. Whilst milder cases are often treated with topical corticosteroids, more severe cases may be  
20 treated with antiproliferative agents, such as the antimetabolite methotrexate, the DNA synthesis inhibitor hydroxyurea, and the microtubule disrupter colchicine.

Other diseases associated with an abnormally high level of cellular proliferation include restenosis, where vascular smooth muscle cells are involved; inflammatory disease states, where endothelial cells, inflammatory cells and glomerular cells are involved; myocardial  
25 infarction, where heart muscle cells are involved; glomerular nephritis, where kidney cells are involved; transplant rejection, where endothelial cells are involved; and infectious diseases such as HIV infection and malaria, where certain immune cells and/or other infected cells are involved.

Inhibition of cellular proliferation can be brought about by several mechanisms,  
30 including; alkylating agents; topoisomerase inhibitors; nucleotide analogues; antibiotics; hormone antagonists; and nucleic acid damaging agents; *inter alia*. One pharmacologically important mechanism of inhibiting cellular proliferation is by means of

## 2.

binding tubulin. Tubulin is an asymmetric dimer composed of alpha and beta subunits, that polymerizes to form structural components of the cytoskeleton called microtubules. Microtubules must be highly dynamic in order to carry out many of their functions. At certain stages of the cell cycle, or in particular cell types or organelles, stable microtubules are required, such as for transport within axons or for ciliary and flagellar movement. Micro-tubules assemble during the G2 phase of the cell cycle, and participate in the formation of the mitotic spindle which facilitates the segregation of sister chromatids during the process of cell division. The essential role of microtubules in cell division and the ability of drugs that interact with tubulin to interfere with the cell cycle have made tubulin a successful target for applications that include anti-cancer drugs, fungicides, and herbicides. Typical tubulin ligands such as colchicine, paclitaxel, the *Vinca* alkaloids such as vinblastine, the epothilones, the halicondrins, benomyl and mebendazole directly inhibit cell division by binding to tubulin which leads to the arrest of the cell cycle at the G2/M boundary of mitosis. This mechanism is the basis of the therapeutic value of compounds of this type, such as treating gout with colchicine, restenosis with paclitaxel, cancer with paclitaxel, vinblastine, epothilones and halichondrins, and fungal infections with benomyl and malaria and helminths with mebendazole.

Interfering with microtubule dynamics or stability can inhibit cell division in several ways. Both stabilizing microtubules or inhibiting their polymerization will prevent the cytoskeleton restructuring that is required at several points in the cell cycle and lead to an arrest of the cell's progression from one stage in the cell cycle to the next. Three main classes of tubulin-binding drugs, namely colchicine analogues, *Vinca* alkaloids, and the taxanes, have been identified, each of which possesses a specific binding site on the  $\beta$ -tubulin molecule. Paclitaxel (Taxol<sup>TM</sup>) and related taxanes represent a class of drugs that stabilize microtubules, a process that ultimately leads to the "freezing" of the microtubule structures so that they cannot be restructured (Jordan M A. and Wilson L., 1998). Subsequent arrest at mitosis induces the apoptotic mechanism to cause cell death. A number of colchicine analogues, as well as several other compounds that bind to the same site on  $\beta$ -tubulin as colchicine disrupt tubulin polymerization and disrupt microtubular formation. Vinblastine and several other vinca-related drugs bind to a site that is distinct from the colchicine site. Compounds that bind at the Vinca-site prevent microtubule formation and destabilize microtubules (Jordan et al, 1986; Rai and Wolff (1996). This invention is therefore directed to compounds that potentially modulate microtubule dynamics by binding to tubulin.

## 3.

Accordingly, the present invention aims to provide compounds which are directly or indirectly toxic to actively dividing cells and are useful in the treatment of cancer, viral and bacterial infections, vascular restenosis, inflammatory diseases, autoimmune diseases, or psoriasis. The present invention is also directed to therapeutic compositions for treating said conditions. Further aspects of the invention are to provide methods for killing actively proliferating cells, such as cancerous, bacterial, or epithelial cells, and treating all types of cancers, infections, inflammatory, and generally proliferative conditions. A further aspect relates to provide methods for treating other medical conditions characterized by the presence of rapidly proliferating cells, such as psoriasis and other skin disorders.

In one embodiment, the method of the invention is used in the treatment of sarcomas, carcinomas and/or leukemias. Exemplary disorders for which the subject method can be used alone or as part of a treatment regimen include: fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms' tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma, and retinoblastoma.

In certain embodiments, the method of the invention is to be used to treat disorders such as carcinomas forming from tissue of the breast, prostate, kidney, bladder or colon.

In other embodiments, the method of the invention is used to treat hyperplastic or neoplastic disorders arising in adipose tissue, such as adipose cell tumors, e.g., lipomas, fibrolipomas, lipoblastomas, lipomatosis, hibernomas, hemangiomas and/or liposarcomas.

In still other embodiments, infectious and parasitic agents (e.g. bacteria, trypanosomes, fungi, etc) can also be controlled using the subject compositions and compounds.




4.

## SUMMARY OF THE INVENTION

The present inventors have found that a group of compounds based upon a substituted heterocyclic scaffold inhibit the growth and proliferation of cancer cells. The present inventors have further shown that these compounds can bind to tubulin. Such compounds would also be useful in the treatment of other hyperproliferation related disorders.

Accordingly, in a first aspect the present invention provides a method of modulating microtubule polymerisation in a subject, the method comprising administering a therapeutically effective amount of at least one compound of the general formula



I

or pharmaceutically acceptable prodrugs, salts, hydrates, solvates, crystal forms or diastereomers thereof, wherein:

R1 is H, C<sub>1-4</sub> alkyl;

15 Q is a bond, or C<sub>1-4</sub> alkyl;

A is aryl, hetaryl optionally substituted with 0-3 substituents independently chosen from halogen, C<sub>1-4</sub> alkyl, CH<sub>2</sub>F, CHF<sub>2</sub>, CF<sub>3</sub>, CN, aryl, hetaryl, OCF<sub>3</sub>, OC<sub>1-4</sub>alkyl, OC<sub>2-5</sub>alkylNR<sub>4</sub>R<sub>5</sub>, Oaryl, Ohetaryl, CO<sub>2</sub>R<sub>4</sub>, CONR<sub>4</sub>R<sub>5</sub>, nitro, NR<sub>4</sub>R<sub>5</sub>, C<sub>1-4</sub> alkylNR<sub>4</sub>R<sub>5</sub>, NR<sub>6</sub>C<sub>1-4</sub>alkylNR<sub>4</sub>R<sub>5</sub>, NR<sub>4</sub>COR<sub>5</sub>, NR<sub>6</sub>CONR<sub>4</sub>R<sub>5</sub>, NR<sub>4</sub>SO<sub>2</sub>R<sub>5</sub>;

20 R<sub>4</sub>, R<sub>5</sub> are each independently H, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl cycloalkyl, C<sub>1-4</sub> alkyl cyclohetalkyl, aryl, hetaryl, C<sub>1-4</sub>alkyl aryl, C<sub>1-4</sub> alkyl hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>7</sub>;

R<sub>6</sub> is selected from H, C<sub>1-4</sub> alkyl;

5.

R7 is selected from H, C<sub>1-4</sub> alkyl, aryl, hetaryl, C<sub>1-4</sub> alkyl aryl, C<sub>1-4</sub> alkyl hetaryl;

5 R2 is 0-2 substituents independently selected from halogen, C<sub>1-4</sub>alkyl, OH, OC<sub>1-4</sub>alkyl, CH<sub>2</sub>F, CHF<sub>2</sub>, CF<sub>3</sub>, OCF<sub>3</sub>, CN, C<sub>1-4</sub>alkylNR<sub>8</sub>R<sub>9</sub>, OC<sub>1-4</sub>alkylNR<sub>8</sub>R<sub>9</sub>, CO<sub>2</sub>R<sub>8</sub>, CONR<sub>8</sub>R<sub>9</sub>, NR<sub>8</sub>R<sub>9</sub>, NR<sub>8</sub>COR<sub>9</sub>, NR<sub>10</sub>CONR<sub>8</sub>R<sub>9</sub>, NR<sub>8</sub>SO<sub>2</sub>R<sub>9</sub>;

R8, R9 are each independently H, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl cycloalkyl, C<sub>1-4</sub> alkyl cyclohetalkyl, aryl, hetaryl, C<sub>1-4</sub> alkyl aryl, C<sub>1-4</sub> alkyl hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>11</sub>;

10 R10 is selected from H, C<sub>1-4</sub> alkyl, aryl or hetaryl;

R11 is selected from H, C<sub>1-4</sub> alkyl, aryl, hetaryl, C<sub>1-4</sub> alkyl aryl, C<sub>1-4</sub> alkyl hetaryl;

Y is halogen, OH, NR<sub>12</sub>R<sub>13</sub>, NR<sub>14</sub>COR<sub>12</sub>, NR<sub>14</sub>CONR<sub>12</sub>R<sub>13</sub>, N<sub>14</sub>SO<sub>2</sub>R<sub>13</sub>;

15 R12 and R13 are each independently H, CH<sub>2</sub>F, CHF<sub>2</sub>, CF<sub>3</sub>, CN, C<sub>1-4</sub> alkyl optionally substituted with OH, OC<sub>1-4</sub>alkyl or NR<sub>15</sub>R<sub>16</sub>, cycloalkyl; cyclohetalkyl, C<sub>1-4</sub> alkyl cycloalkyl, C<sub>1-4</sub> alkyl cyclohetalkyl, or may be joined to form an optionally substituted 3-6 membered ring optionally containing an atom selected from O, S, NR<sub>14</sub>

R14, R15 and R16 are each independently selected from H, C<sub>1-4</sub> alkyl;

20 n = 0-4;


W is selected from H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl; where C<sub>1-4</sub>alkyl or C<sub>2-6</sub>alkenyl may be optionally substituted with C<sub>1-4</sub>alkyl, OH, OC<sub>1-4</sub>alkyl, NR<sub>15</sub>R<sub>16</sub>;

25 R15, and R16 are each independently H, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl cycloalkyl, C<sub>1-4</sub> alkyl cyclohetalkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>17</sub>;

R17 is selected from H, C<sub>1-4</sub> alkyl.

In a second aspect the present invention provides a compound of the general formula

6.



II

or pharmaceutically acceptable prodrugs, salts, hydrates, solvates, crystal forms or diastereomers thereof, wherein:

5  $R1$  is H,  $C_{1-4}$  alkyl;

$Q$  is a bond, or  $C_{1-4}$  alkyl;

$A$  is aryl, hetaryl optionally substituted with 0-3 substituents independently chosen from halogen,  $C_{1-4}$  alkyl,  $CH_2F$ ,  $CHF_2$ ,  $CF_3$ , CN, aryl, hetaryl,  $OCF_3$ ,  $OC_{1-4}alkyl$ ,  $OC_{2-5}alkylNR_4R_5$ ,  $Oaryl$ ,  $Ohetaryl$ ,  $CO_2R_4$ ,  $CONR_4R_5$ , nitro,  $NR_4R_5$ ,  $C_{1-4}alkylNR_4R_5$ ,  $NR_6C_{1-4}alkylNR_4R_5$ ,  $NR_4COR_5$ ,  $NR_6CONR_4R_5$ ,  $NR_4SO_2R_5$ ;

$R_4$ ,  $R_5$  are each independently H,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkyl cycloalkyl,  $C_{1-4}$  alkyl cyclohetalkyl, aryl, hetaryl,  $C_{1-4}alkyl$  aryl,  $C_{1-4}alkyl$  hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S,  $NR_7$ ;

15  $R_6$  is selected from H,  $C_{1-4}$  alkyl;


$R_7$  is selected from H,  $C_{1-4}$  alkyl, aryl, hetaryl,  $C_{1-4}alkyl$  aryl,  $C_{1-4}alkyl$  hetaryl;

$R_2$  is 0-2 substituents independently selected from  $C_{1-4}alkyl$  and  $OC_{1-4}alkyl$ ;


20  $Y$  is  $CH_2OH$ ,  $OC_{1-4}alkylOH$ ,  $OC_{1-4}alkylR_{12}$ ,  $OC_{1-4}alkylNR_{12}NR_{13}$ ,  $C(O)R_{12}$ ,  $CH_2R_{12}$ ,  $COOR_{12}$ ,  $CONR_{12}R_{13}$ ,  $CONR_{12}R_{13}$ ,  $CH_2NR_{12}R_{13}$ ,  $NHCO R_{12}$ ,  $NHCONR_{12}R_{13}$ ,

$R_{12}$  and  $R_{13}$  are each independently H,  $C_{1-2}$  alkyl,  $(CH_2)_3NEt_2$ ,  $(CH_2)_2NMe_2$ ,  $(CH_2)_5NH_2$ ,  $(CH_2)_2OH$ ,


7.




5



10



15



n = 0-4;

W is selected from H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl; where C<sub>1-4</sub>alkyl or C<sub>2-6</sub>alkenyl may be optionally substituted with C<sub>1-4</sub>alkyl, OH, OC<sub>1-4</sub>alkyl, NR<sub>15</sub>R<sub>16</sub>;

20

R<sub>15</sub>, and R<sub>16</sub> are each independently H, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl cycloalkyl, C<sub>1-4</sub> alkyl cyclohetalkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>17</sub>

R<sub>17</sub> is selected from H, C<sub>1-4</sub> alkyl;




8.

wherein when Y is CH<sub>2</sub>R<sub>12</sub> then R<sub>12</sub> is not H, C<sub>1-2</sub>alkyl.

In a third aspect, the present invention provides a composition comprising a carrier and at least one compound of the second aspect.

In a fourth aspect the present invention provides a method of treating a  
 5 hyperproliferation-related disorder in a subject, the method comprising administering a therapeutically effective amount of at least one compound of the second aspect or a composition of the third aspect.

In a fifth aspect the present invention provides a compound of the general formula



III

or pharmaceutically acceptable prodrugs, salts, hydrates, solvates, crystal forms or diastereomers thereof, wherein:

15 X<sub>1</sub>, X<sub>2</sub>, X<sub>3</sub>, X<sub>4</sub> are selected from the following:

- (i) X<sub>1</sub> and X<sub>2</sub> are N and X<sub>3</sub> and X<sub>4</sub> are C independently substituted with Y;
- (ii) X<sub>1</sub> and X<sub>4</sub> are N and X<sub>2</sub> and X<sub>3</sub> are C independently substituted with Y;
- (iii) X<sub>1</sub> and X<sub>3</sub> are N and X<sub>2</sub> and X<sub>4</sub> are C independently substituted with Y;
- (iv) X<sub>2</sub> and X<sub>4</sub> are N and X<sub>1</sub> and X<sub>3</sub> are C independently substituted with Y;
- 20 (v) X<sub>1</sub> is N and X<sub>2</sub>, X<sub>3</sub>, and X<sub>4</sub> are C independently substituted with Y;

9.

- (vi)  $X_3$  is N and  $X_1$ ,  $X_2$ , and  $X_4$  are C independently substituted with Y;  
(vii)  $X_4$  is N and  $X_1$ ,  $X_2$ , and  $X_3$  are C independently substituted with Y;  
(viii)  $X_2$  is N and  $X_1$ ,  $X_3$ , and  $X_4$  are C independently substituted with Y; and  
(ix)  $X_1$ ,  $X_2$  and  $X_3$  are N and  $X_4$  is C substituted with Y;

5 R1 is H,  $C_{1-6}$ alkyl,  $C_{1-6}$  alkylNR5R6,  $C_{1-6}$ alkylNR5COR6,  $C_{1-6}$ alkylNR5SO<sub>2</sub>R6,  $C_{1-6}$ alkylCO<sub>2</sub>R5,  $C_{1-6}$ alkylCONR5R6;

R5 and R6 are each independently H,  $C_{1-4}$ alkyl, aryl, hetaryl,  $C_{1-4}$ alkylaryl,  $C_{1-4}$ alkylhetaryl or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR7;

10 R7 is selected from H,  $C_{1-4}$ alkyl;

R2 is selected from  $C_{1-6}$ alkylOH, OC<sub>2-6</sub>alkylOH,  $C_{1-6}$ alkylNR8R9, OC<sub>2-6</sub>alkylNR8R9,  $C_{1-6}$ alkylNR8COR9, OC<sub>2-6</sub>alkylNR8COR9,  $C_{1-6}$ alkylhetaryl, OC<sub>2-6</sub>alkylhetaryl, OCONR8R9, NR8COOR9, NR10CONR8R9, CONR8R9, NR8COR12;

15 R8, R9 are each independently H,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkylNR11R13, hetaryl, cyclohetalkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR14;

R12 is  $C_{2-4}$ alkyl,  $C_{1-4}$ alkylNR11R13, hetaryl, cyclohetalkyl;

20 R11, R13 are each independently H,  $C_{1-4}$ alkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR14;

R14 is selected from H,  $C_{1-4}$ alkyl;

R10 is H,  $C_{1-4}$  alkyl;

R3 and R4 are each independently H, halogen,  $C_{1-4}$ alkyl, OH, OC<sub>1-4</sub>alkyl, CF<sub>3</sub>, OCF<sub>3</sub>;

25 Q is a bond, or  $C_{1-4}$  alkyl;

W is selected from H,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl; where  $C_{1-4}$ alkyl or  $C_{2-6}$ alkenyl may be optionally substituted with  $C_{1-4}$ alkyl, OH, OC<sub>1-4</sub>alkyl, NR15R16;

10.

R15, and R16 are each independently H, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyl cycloalkyl, C<sub>1-4</sub>alkyl cyclohetalkyl, aryl, hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR17;

5 R17 is selected from H, C<sub>1-4</sub>alkyl;

A is aryl, hetaryl optionally substituted with 0-3 substituents independently chosen from halogen, C<sub>1-4</sub> alkyl, CF<sub>3</sub>, aryl, hetaryl, OCF<sub>3</sub>, OC<sub>1-4</sub>alkyl, OC<sub>2-5</sub>alkylNR18R19, Oaryl, Ohetaryl, CO<sub>2</sub>R18, CONR18R19, NR18R19, C<sub>1-4</sub>alkylNR18R19, NR20C<sub>1-4</sub>alkylNR18R19, NR18COR19, NR20CONR18R19, NR18SO<sub>2</sub>R19;

R18, R19 are each independently H, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl cyclohetalkyl, aryl, hetaryl, C<sub>1-4</sub>alkyl aryl, C<sub>1-4</sub> alkyl hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR21;

15 R21 is selected from H, C<sub>1-4</sub>alkyl;

R20 is selected from H, C<sub>1-4</sub>alkyl;

Y is selected from H, C<sub>1-4</sub>alkyl, OH, NR22R23;

R22, R23 are each independently H, C<sub>1-4</sub>alkyl.


20 In a sixth aspect, the present invention provides a composition comprising a carrier and one or more compounds of the fifth aspect.

In a seventh aspect, the present invention provides a method of treatment of a hyperproliferation-related disorder in a subject, the method comprising administering a therapeutically effective amount of at least one of the compounds of the fifth aspect or a composition according to the sixth aspect.

25 In an eighth aspect, the present invention consists in a method of treating a protein-kinase related disorder in a subject, the method comprising administering a therapeutically effective amount of at least one of the compounds of the fifth aspect or a composition according to the sixth aspect.

In a ninth aspect, the present invention provides a compound of the general formula

11.



V

or pharmaceutically acceptable prodrugs, salts, hydrates, solvates, crystal forms or  
 5 diastereomers thereof, wherein:

$X_1, X_2, X_3, X_4$  are selected from the following:

- (i)  $X_1$  and  $X_2$  are N and  $X_3$  and  $X_4$  are C independently substituted with Y;
- (ii)  $X_1$  and  $X_4$  are N and  $X_2$  and  $X_3$  are C independently substituted with Y;
- (iii)  $X_2$  and  $X_4$  are N and  $X_1$  and  $X_3$  are C independently substituted with Y;
- 10 (iv)  $X_1$  is N and  $X_2, X_3$ , and  $X_4$  are C independently substituted with Y;
- (v)  $X_3$  is N and  $X_1, X_2$ , and  $X_4$  are C independently substituted with Y;
- (vi)  $X_4$  is N and  $X_1, X_2$ , and  $X_3$  are C independently substituted with Y;
- (vii)  $X_2$  is N and  $X_1, X_3$ , and  $X_4$  are C independently substituted with Y; and
- (viii)  $X_1, X_2$  and  $X_3$  are N and  $X_4$  is C substituted with Y;

15  $R_1$  is H,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkylNR<sub>5</sub>R<sub>6</sub>,  $C_{1-6}$ alkylNR<sub>5</sub>COR<sub>6</sub>,  $C_{1-6}$ alkylNR<sub>5</sub>SO<sub>2</sub>R<sub>6</sub>,  
 $C_{1-6}$ alkylCO<sub>2</sub>R<sub>5</sub>,  $C_{1-6}$ alkylCONR<sub>5</sub>R<sub>6</sub>;

$R_5$  and  $R_6$  are each independently H,  $C_{1-4}$ alkyl, aryl, hetaryl,  $C_{1-4}$ alkylaryl,  
 $C_{1-4}$ alkylhetaryl or may be joined to form an optionally substituted 3-8  
 membered ring optionally containing an atom selected from O, S, NR<sub>7</sub>;

20  $R_7$  is selected from H,  $C_{1-4}$ alkyl;

12.

R2 is selected from OH, OC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylOH, OC<sub>2-6</sub>alkylOH, C<sub>1-6</sub>alkylNR<sub>8</sub>R<sub>9</sub>, OC<sub>2-6</sub>alkylNR<sub>8</sub>R<sub>9</sub>, C<sub>1-6</sub>alkylNR<sub>8</sub>COR<sub>9</sub>, OC<sub>2-6</sub>alkylNR<sub>8</sub>COR<sub>9</sub>, C<sub>1-6</sub>alkylhetaryl, OC<sub>2-6</sub>alkylhetaryl, OCONR<sub>8</sub>R<sub>9</sub>, NR<sub>8</sub>COOR<sub>9</sub>, NR<sub>10</sub>CONR<sub>8</sub>R<sub>9</sub>, CONR<sub>8</sub>R<sub>9</sub>, NR<sub>8</sub>COR<sub>12</sub>;

5 R<sub>8</sub>, R<sub>9</sub> are each independently H, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylNR<sub>11</sub>R<sub>13</sub>, hetaryl, cyclohetalkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>14</sub>;

R<sub>12</sub> is C<sub>2-4</sub>alkyl, C<sub>1-4</sub>alkylNR<sub>11</sub>R<sub>13</sub>, hetaryl, cyclohetalkyl;

10 R<sub>11</sub>, R<sub>13</sub> are each independently H, C<sub>1-4</sub>alkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>14</sub>;

R<sub>14</sub> is selected from H, C<sub>1-4</sub>alkyl;

R<sub>10</sub> is H, C<sub>1-4</sub> alkyl;

15 R<sub>3</sub> and R<sub>4</sub> are each independently H, halogen, C<sub>1-4</sub>alkyl, OH, OC<sub>1-4</sub>alkyl, CF<sub>3</sub>, OCF<sub>3</sub>;

Q is a bond, or C<sub>1-4</sub>alkyl;

W is selected from H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl; where C<sub>1-4</sub>alkyl or C<sub>2-6</sub>alkenyl may be optionally substituted with C<sub>1-4</sub>alkyl, OH, OC<sub>1-4</sub>alkyl, NR<sub>15</sub>R<sub>16</sub>;

20 R<sub>15</sub>, and R<sub>16</sub> are each independently H, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyl cycloalkyl, C<sub>1-4</sub>alkyl cyclohetalkyl, aryl, hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>17</sub>;

R<sub>17</sub> is selected from H, C<sub>1-4</sub>alkyl;

25 A is aryl, hetaryl optionally substituted with 0-3 substituents independently chosen from halogen, C<sub>1-4</sub> alkyl, CF<sub>3</sub>, aryl, hetaryl, OCF<sub>3</sub>, OC<sub>1-4</sub>alkyl, OC<sub>2-5</sub>alkylNR<sub>18</sub>R<sub>19</sub>, Oaryl, Ohetaryl, CO<sub>2</sub>R<sub>18</sub>, CONR<sub>18</sub>R<sub>19</sub>, NR<sub>18</sub>R<sub>19</sub>, C<sub>1-4</sub>alkylNR<sub>18</sub>R<sub>19</sub>, NR<sub>20</sub>C<sub>1-4</sub>alkylNR<sub>18</sub>R<sub>19</sub>, NR<sub>18</sub>COR<sub>19</sub>, NR<sub>20</sub>CONR<sub>18</sub>R<sub>19</sub>, NR<sub>18</sub>SO<sub>2</sub>R<sub>19</sub>;



13.

R18, R19 are each independently H, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl cyclohetalkyl, aryl, hetaryl, C<sub>1-4</sub>alkyl aryl, C<sub>1-4</sub> alkyl hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR21;

5 R21 is selected from H, C<sub>1-4</sub> alkyl;

R20 is selected from H, C<sub>1-4</sub> alkyl;

Y is selected from H, C<sub>1-4</sub>alkyl, OH, NR22R23;

R22, R23 are each independently H, C<sub>1-4</sub> alkyl.


10 In a tenth aspect, the present invention provides a composition comprising a carrier and one or more compounds of the ninth aspect.

In an eleventh aspect, the present invention provides a method of treatment of a hyperproliferation-related disorder in a subject, the method comprising administering a therapeutically effective amount of at least one of the compounds of the ninth aspect or a composition according to the tenth aspect.

15 In a twelfth aspect, the present invention consists in a method of treating a protein-kinase related disorder in a subject, the method comprising administering a therapeutically effective amount of at least one of the compounds of the ninth aspect or a composition according to the tenth aspect.

## DETAILED DESCRIPTION OF THE INVENTION

20 In a first aspect the present invention provides a method of modulating microtubule polymerisation in a subject, the method comprising administering a therapeutically effective amount of at least one compound of the general formula



I

14.

or pharmaceutically acceptable salts, hydrates, solvates, crystal forms or diastereomers thereof, wherein:

R1 is H, C<sub>1-4</sub> alkyl;

Q is a bond, or C<sub>1-4</sub> alkyl;

- 5 A is aryl, hetaryl optionally substituted with 0-3 substituents independently chosen from halogen, C<sub>1-4</sub> alkyl, CH<sub>2</sub>F, CHF<sub>2</sub>, CF<sub>3</sub>, CN, aryl, hetaryl, OCF<sub>3</sub>, OC<sub>1-4</sub>alkyl, OC<sub>2-5</sub>alkylNR<sub>4</sub>R<sub>5</sub>, Oaryl, Ohetaryl, CO<sub>2</sub>R<sub>4</sub>, CONR<sub>4</sub>R<sub>5</sub>, nitro, NR<sub>4</sub>R<sub>5</sub>, C<sub>1-4</sub> alkylNR<sub>4</sub>R<sub>5</sub>, NR<sub>6</sub>C<sub>1-4</sub>alkylNR<sub>4</sub>R<sub>5</sub>, NR<sub>4</sub>COR<sub>5</sub>, NR<sub>6</sub>CONR<sub>4</sub>R<sub>5</sub>, NR<sub>4</sub>SO<sub>2</sub>R<sub>5</sub>; and R<sub>4</sub>, R<sub>5</sub> are each independently H, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl cycloalkyl, C<sub>1-4</sub> alkyl cyclohetalkyl, aryl, hetaryl, C<sub>1-4</sub>alkyl aryl, C<sub>1-4</sub> alkyl hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>7</sub>; and R<sub>6</sub> is selected from H, C<sub>1-4</sub> alkyl; and R<sub>7</sub> is selected from H, C<sub>1-4</sub> alkyl, aryl, hetaryl, C<sub>1-4</sub> alkyl aryl, C<sub>1-4</sub> alkyl hetaryl;
- 10
- 15 R<sub>2</sub> is 0-2 substituents independently selected from halogen, C<sub>1-4</sub>alkyl, OH, OC<sub>1-4</sub>alkyl, CH<sub>2</sub>F, CHF<sub>2</sub>, CF<sub>3</sub>, OCF<sub>3</sub>, CN, C<sub>1-4</sub>alkylNR<sub>8</sub>R<sub>9</sub>, OC<sub>1-4</sub>alkylNR<sub>8</sub>R<sub>9</sub>, CO<sub>2</sub>R<sub>8</sub>, CONR<sub>8</sub>R<sub>9</sub>, NR<sub>8</sub>R<sub>9</sub>, NR<sub>8</sub>COR<sub>9</sub>, NR<sub>10</sub>CONR<sub>8</sub>R<sub>9</sub>, NR<sub>8</sub>SO<sub>2</sub>R<sub>9</sub>; and R<sub>8</sub>, R<sub>9</sub> are each independently H, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl cycloalkyl, C<sub>1-4</sub> alkyl cyclohetalkyl, aryl, hetaryl, C<sub>1-4</sub> alkyl aryl, C<sub>1-4</sub> alkyl hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>11</sub>; and R<sub>10</sub> is selected from H, C<sub>1-4</sub> alkyl, aryl or hetaryl; and R<sub>11</sub> is selected from H, C<sub>1-4</sub> alkyl, aryl, hetaryl, C<sub>1-4</sub> alkyl aryl, C<sub>1-4</sub> alkyl hetaryl;
- 20
- 25 Y is halogen; OH; NR<sub>12</sub>R<sub>13</sub>; NR<sub>14</sub>COR<sub>12</sub>; NR<sub>14</sub>CONR<sub>12</sub>R<sub>13</sub>; NR<sub>14</sub>SO<sub>2</sub>R<sub>13</sub>; and R<sub>12</sub> and R<sub>13</sub> are each independently H; CH<sub>2</sub>F; CHF<sub>2</sub>; CF<sub>3</sub>; CN; C<sub>1-4</sub> alkyl optionally substituted with OH, OC<sub>1-4</sub>alkyl or NR<sub>15</sub>R<sub>16</sub>; cycloalkyl; cyclohetalkyl; C<sub>1-4</sub> alkyl cycloalkyl; C<sub>1-4</sub> alkyl cyclohetalkyl; or may be joined to form an optionally substituted 3-6 membered ring optionally containing an atom selected from O, S, NR<sub>14</sub> and R<sub>14</sub>, R<sub>15</sub> and R<sub>16</sub> are each independently selected from H, C<sub>1-4</sub> alkyl;
- 30 n = 0-4;
- W is selected from H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl; where C<sub>1-4</sub>alkyl or C<sub>2-6</sub>alkenyl may be optionally substituted with C<sub>1-4</sub>alkyl, OH, OC<sub>1-4</sub>alkyl, NR<sub>15</sub>R<sub>16</sub>; and R<sub>15</sub>, and R<sub>16</sub> are each independently H, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl cycloalkyl, C<sub>1-4</sub> alkyl cyclohetalkyl,


15.

or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR17 and R17 is selected from H, C<sub>1-4</sub> alkyl.

Preferably, the method is used in the treatment of a hyperproliferation-related disorder.

Preferably, the compound is selected from the compounds set out in Table 2.

5 In a second aspect the present invention provides a compound of the general formula



II

or pharmaceutically acceptable prodrugs, salts, hydrates, solvates, crystal forms or diastereomers thereof, wherein:

10 R1 is H, C<sub>1-4</sub> alkyl;


Q is a bond, or C<sub>1-4</sub> alkyl;

A is aryl, hetaryl optionally substituted with 0-3 substituents independently chosen from halogen, C<sub>1-4</sub> alkyl, CH<sub>2</sub>F, CHF<sub>2</sub>, CF<sub>3</sub>, CN, aryl, hetaryl, OCF<sub>3</sub>, OC<sub>1-4</sub>alkyl, OC<sub>2-5</sub>alkylNR<sub>4</sub>R<sub>5</sub>, Oaryl, Ohetaryl, CO<sub>2</sub>R<sub>4</sub>, CONR<sub>4</sub>R<sub>5</sub>, nitro, NR<sub>4</sub>R<sub>5</sub>, C<sub>1-4</sub> alkylNR<sub>4</sub>R<sub>5</sub>, NR<sub>6</sub>C<sub>1-4</sub>alkylNR<sub>4</sub>R<sub>5</sub>, NR<sub>4</sub>COR<sub>5</sub>, NR<sub>6</sub>CONR<sub>4</sub>R<sub>5</sub>, NR<sub>4</sub>SO<sub>2</sub>R<sub>5</sub>; and  
 15 R<sub>4</sub>, R<sub>5</sub> are each independently H, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl cycloalkyl, C<sub>1-4</sub> alkyl cyclohetalkyl, aryl, hetaryl, C<sub>1-4</sub>alkyl aryl, C<sub>1-4</sub> alkyl hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>7</sub>; and R<sub>6</sub> is selected from H, C<sub>1-4</sub> alkyl; and R<sub>7</sub> is selected  
 20 from H, C<sub>1-4</sub> alkyl, aryl, hetaryl, C<sub>1-4</sub> alkyl aryl, C<sub>1-4</sub> alkyl hetaryl;

R<sub>2</sub> is 0-2 substituents independently selected from C<sub>1-4</sub>alkyl and OC<sub>1-4</sub>alkyl;

Y is CH<sub>2</sub>OH, OC<sub>1-4</sub>alkylOH, OC<sub>1-4</sub>alkylR<sub>12</sub>, OC<sub>1-4</sub>alkylNR<sub>12</sub>NR<sub>13</sub>, C(O)R<sub>12</sub>, CH<sub>2</sub>R<sub>12</sub>, COOR<sub>12</sub>, CONR<sub>12</sub>R<sub>13</sub>, OCONR<sub>12</sub>R<sub>13</sub>, CH<sub>2</sub>NR<sub>12</sub>R<sub>13</sub>, NHCOR<sub>12</sub>, NHCONR<sub>12</sub>R<sub>13</sub>, and R<sub>12</sub> and R<sub>13</sub> are each independently H, C<sub>1-2</sub> alkyl,  
 25 (CH<sub>2</sub>)<sub>3</sub>NEt<sub>2</sub>, (CH<sub>2</sub>)<sub>2</sub>NMe<sub>2</sub>, (CH<sub>2</sub>)<sub>5</sub>NH<sub>2</sub>, (CH<sub>2</sub>)<sub>2</sub>OH,

16.



20 W is selected from H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl; where C<sub>1-4</sub>alkyl or C<sub>2-6</sub>alkenyl may be optionally substituted with C<sub>1-4</sub>alkyl, OH, OC<sub>1-4</sub>alkyl, NR<sub>15</sub>R<sub>16</sub>; and R<sub>15</sub>, and R<sub>16</sub> are each independently H, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl cycloalkyl, C<sub>1-4</sub> alkyl cyclohetalkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>17</sub> and R<sub>17</sub> is selected from H, C<sub>1-4</sub> alkyl;

wherein when Y is CH<sub>2</sub>R<sub>12</sub> then R<sub>12</sub> is not H, C<sub>1-2</sub>alkyl.

17.


In a third aspect, the present invention comprises a composition comprising a carrier and at least one compound of the second aspect.

In a fourth aspect the present invention comprises a method of treating a hyperproliferation-related disorder in a subject, the method comprising administering a therapeutically effective amount of at least one compound of the second aspect or a composition of the third aspect.

Preferably, the hyperproliferation-related disorder is treatable by the modulation of microtubule polymerisation.

In a fifth aspect the present invention provides a compound of the general formula

10



III

or pharmaceutically acceptable prodrugs, salts, hydrates, solvates, crystal forms or diastereomers thereof, wherein:

15

$X_1, X_2, X_3, X_4$  are selected from the following:

- (i)  $X_1$  and  $X_2$  are N and  $X_3$  and  $X_4$  are C independently substituted with Y;
- (ii)  $X_1$  and  $X_4$  are N and  $X_2$  and  $X_3$  are C independently substituted with Y;
- (iii)  $X_1$  and  $X_3$  are N and  $X_2$  and  $X_4$  are C independently substituted with Y;
- (iv)  $X_2$  and  $X_4$  are N and  $X_1$  and  $X_3$  are C independently substituted with Y;
- (v)  $X_1$  is N and  $X_2, X_3$ , and  $X_4$  are C independently substituted with Y;

20



18.

- (vi)  $X_3$  is N and  $X_1$ ,  $X_2$ , and  $X_4$  are C independently substituted with Y;
- (vii)  $X_4$  is N and  $X_1$ ,  $X_2$ , and  $X_3$  are C independently substituted with Y;
- (viii)  $X_2$  is N and  $X_1$ ,  $X_3$ , and  $X_4$  are C independently substituted with Y; and
- (ix)  $X_1$ ,  $X_2$  and  $X_3$  are N and  $X_4$  is C substituted with Y;

5 R1 is H,  $C_{1-6}$ alkyl,  $C_{1-6}$  alkylNR5R6,  $C_{1-6}$ alkylNR5COR6,  $C_{1-6}$ alkylNR5SO<sub>2</sub>R6,  $C_{1-6}$ alkylCO<sub>2</sub>R5,  $C_{1-6}$ alkylCONR5R6, where R5 and R6 are each independently H,  $C_{1-4}$ alkyl, aryl, hetaryl,  $C_{1-4}$ alkylaryl,  $C_{1-4}$ alkylhetaryl or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR7;

10 R7 is selected from H,  $C_{1-4}$ alkyl;

R2 is selected from  $C_{1-6}$ alkylOH, OC<sub>2-6</sub>alkylOH,  $C_{1-6}$ alkylNR8R9, OC<sub>2-6</sub>alkylNR8R9,  $C_{1-6}$ alkylNR8COR9, OC<sub>2-6</sub>alkylNR8COR9,  $C_{1-6}$ alkylhetaryl, OC<sub>2-6</sub>alkylhetaryl, OCONR8R9, NR8COOR9, NR10CONR8R9, CONR8R9, NR8COR12;

15 R8, R9 are each independently H,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkylNR11R13, hetaryl, cyclohetalkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR14;

R12 is  $C_{2-4}$ alkyl,  $C_{1-4}$ alkylNR11R13, hetaryl, cyclohetalkyl;

20 R11, R13 are each independently H,  $C_{1-4}$ alkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR14;

R14 is selected from H,  $C_{1-4}$ alkyl;

R10 is H,  $C_{1-4}$  alkyl;

R3 and R4 are each independently H, halogen,  $C_{1-4}$ alkyl, OH, OC<sub>1-4</sub>alkyl, CF<sub>3</sub>, OCF<sub>3</sub>;

25 Q is a bond, or  $C_{1-4}$  alkyl;

W is selected from H,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl; where  $C_{1-4}$ alkyl or  $C_{2-6}$ alkenyl may be optionally substituted with  $C_{1-4}$ alkyl, OH, OC<sub>1-4</sub>alkyl, NR15R16;

19.

R15, and R16 are each independently H, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyl cycloalkyl, C<sub>1-4</sub>alkyl cyclohetalkyl, aryl, hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR17;

5 R17 is selected from H, C<sub>1-4</sub>alkyl;

A is aryl, hetaryl optionally substituted with 0-3 substituents independently chosen from halogen, C<sub>1-4</sub> alkyl, CF<sub>3</sub>, aryl, hetaryl, OCF<sub>3</sub>, OC<sub>1-4</sub>alkyl, OC<sub>2-5</sub>alkylNR18R19, Oaryl, Ohetaryl, CO<sub>2</sub>R18, CONR18R19, NR18R19, C<sub>1-4</sub> alkylNR18R19, NR20C<sub>1-4</sub>alkylNR18R19, NR18COR19, NR20CONR18R19, NR18SO<sub>2</sub>R19;

R18, R19 are each independently H, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl cyclohetalkyl, aryl, hetaryl, C<sub>1-4</sub>alkyl aryl, C<sub>1-4</sub> alkyl hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR21;

15 R21 is selected from H, C<sub>1-4</sub>alkyl;

R20 is selected from H, C<sub>1-4</sub>alkyl;

Y is selected from H, C<sub>1-4</sub>alkyl, OH, NR22R23;

R22, R23 are each independently H, C<sub>1-4</sub>alkyl.

In the above description it will be appreciated that:

20 C<sub>1-4</sub> alkyl means an unsubstituted or optionally substituted straight or branched alkyl chain

Aryl means unsubstituted or optionally substituted phenyl or naphthyl.


Hetaryl means an unsubstituted or optionally substituted 5- or 6-membered heteroaromatic ring containing one or more heteroatoms selected from O, N, S.

25 Cycloalkyl means a 3-8 membered saturated ring

Cyclohetalkyl means an optionally substituted 3-8 membered saturated ring containing 1-3 heteroatoms selected from O, S, NR24, where R24 is H, C<sub>1-4</sub> alkyl, aryl, hetaryl.

20.

In a preferred embodiment the compound is selected from compounds of the general formula



IV

5 or pharmaceutically acceptable prodrugs, salts, hydrates, solvates, crystal forms or diastereomers thereof, wherein:

$X_1, X_2, X_3, X_4$  are selected from the following:

- (i)  $X_1$  and  $X_2$  are N and  $X_3$  and  $X_4$  are C independently substituted with Y;
- (ii)  $X_1$  and  $X_4$  are N and  $X_2$  and  $X_3$  are C independently substituted with Y;
- 10 (iii)  $X_1$  and  $X_3$  are N and  $X_2$  and  $X_4$  are C independently substituted with Y;
- (iv)  $X_2$  and  $X_4$  are N and  $X_1$  and  $X_3$  are C independently substituted with Y;
- (v)  $X_1$  is N and  $X_2, X_3$ , and  $X_4$  are C independently substituted with Y;
- (vi)  $X_3$  is N and  $X_1, X_2$ , and  $X_4$  are C independently substituted with Y;
- (vii)  $X_4$  is N and  $X_1, X_2$ , and  $X_3$  are C independently substituted with Y;
- 15 (viii)  $X_2$  is N and  $X_1, X_3$ , and  $X_4$  are C independently substituted with Y; and
- (ix)  $X_1, X_2$  and  $X_3$  are N and  $X_4$  is C substituted with Y;

$R_1$  is H,  $C_{1-6}$ alkyl,  $C_{1-6}$  alkylNR<sub>5</sub>R<sub>6</sub>, where R<sub>5</sub> and R<sub>6</sub> are each independently H,  $C_{1-4}$ alkyl, aryl, hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>7</sub>;

20  $R_7$  is selected from H,  $C_{1-4}$  alkyl;

21.

R2 is selected from C<sub>1-6</sub>alkylOH, OC<sub>2-6</sub>alkylOH, C<sub>1-6</sub>alkylNR<sub>8</sub>R<sub>9</sub>, OC<sub>2-6</sub>alkylNR<sub>8</sub>R<sub>9</sub>, C<sub>1-6</sub>alkylNR<sub>8</sub>COR<sub>9</sub>, OC<sub>2-6</sub>alkylNR<sub>8</sub>COR<sub>9</sub>, C<sub>1-6</sub>alkylhetaryl, OC<sub>2-6</sub>alkylhetaryl, OCONR<sub>8</sub>R<sub>9</sub>, NR<sub>8</sub>COOR<sub>9</sub>, NR<sub>10</sub>CONR<sub>8</sub>R<sub>9</sub>, CONR<sub>8</sub>R<sub>9</sub>, NR<sub>8</sub>COR<sub>12</sub>;

5 R<sub>8</sub>, R<sub>9</sub> are each independently H, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylNR<sub>11</sub>R<sub>13</sub>, hetaryl, cyclohetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>14</sub>;

R<sub>12</sub> is C<sub>2-4</sub>alkyl, C<sub>1-4</sub>alkylNR<sub>11</sub>R<sub>13</sub>, hetaryl, cyclohetaryl;

10 R<sub>11</sub>, R<sub>13</sub> are each independently H, C<sub>1-4</sub>alkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>14</sub>;

R<sub>14</sub> is selected from H, C<sub>1-4</sub>alkyl;

R<sub>10</sub> is H, C<sub>1-4</sub>alkyl;

R<sub>3</sub> and R<sub>4</sub> are each independently H, halogen, C<sub>1-4</sub>alkyl, OH, OC<sub>1-4</sub>alkyl, CF<sub>3</sub>, OCF<sub>3</sub>;

15 Q is CH;

W is selected from C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl; where C<sub>1-4</sub>alkyl or C<sub>2-6</sub>alkenyl may be optionally substituted with C<sub>1-4</sub>alkyl, OH, OC<sub>1-4</sub>alkyl, NR<sub>15</sub>R<sub>16</sub>;

20 R<sub>15</sub>, and R<sub>16</sub> are each independently H, C<sub>1-4</sub>alkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>17</sub>;

R<sub>17</sub> is selected from H, C<sub>1-4</sub>alkyl;

25 A is aryl, hetaryl optionally substituted with 0-2 substituents independently chosen from halogen, C<sub>1-4</sub>alkyl, CF<sub>3</sub>, aryl, hetaryl, OCF<sub>3</sub>, OC<sub>1-4</sub>alkyl, OC<sub>2-5</sub>alkylNR<sub>18</sub>R<sub>19</sub>, Oaryl, Ohetaryl, CO<sub>2</sub>R<sub>18</sub>, CONR<sub>18</sub>R<sub>19</sub>, NR<sub>18</sub>R<sub>19</sub>, C<sub>1-4</sub>alkylNR<sub>18</sub>R<sub>19</sub>, NR<sub>20</sub>C<sub>1-4</sub>alkylNR<sub>18</sub>R<sub>19</sub>, NR<sub>18</sub>COR<sub>19</sub>, NR<sub>20</sub>CONR<sub>18</sub>R<sub>19</sub>, NR<sub>18</sub>SO<sub>2</sub>R<sub>19</sub>;

R<sub>18</sub>, R<sub>19</sub> are each independently H, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyl cyclohetaryl, aryl, hetaryl, C<sub>1-4</sub>alkyl aryl, C<sub>1-4</sub>alkyl hetaryl, or may be joined to form an

22.

optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR21;

R21 is selected from H, C<sub>1-4</sub>alkyl;

R20 is selected from H, C<sub>1-4</sub>alkyl;

5 Y is selected from H, C<sub>1-4</sub>alkyl, NR22R23;

R22, R23 are each independently H, C<sub>1-4</sub>alkyl.

In the above description it will be appreciated that:

C<sub>1-4</sub> alkyl means an unsubstituted or optionally substituted straight or branched alkyl chain

10 Aryl means unsubstituted or optionally substituted phenyl or naphthyl.

Hetaryl means an unsubstituted or optionally substituted 5- or 6-membered heteroaromatic ring containing one or more heteroatoms selected from O, N, S.

Cycloalkyl means a 3-8 membered saturated ring

15 Cyclohetalkyl means an optionally substituted 3-8 membered saturated ring containing 1-3 heteroatoms selected from O, S, NR24, where R24 is H, C<sub>1-4</sub> alkyl, aryl, hetaryl.

In a sixth aspect, the present invention provides a composition comprising a carrier and one or more compounds of the fifth aspect.

20 In a seventh aspect, the present invention provides a method of treatment of a hyperproliferation-related disorder in a subject, the method comprising administering a therapeutically effective amount of at least one of the compounds of the fifth aspect or a composition according to the sixth aspect.

Preferably, the hyperproliferation-related disorder is treatable by the modulation of microtubule polymerisation.


25 In an eighth aspect, the present invention consists in a method of treating a protein-kinase related disorder in a subject, the method comprising administering a therapeutically



23.

effective amount of at least one of the compounds of the fifth aspect or a composition according to the sixth aspect.

In a ninth aspect, the present invention provides a compound of the general formula



V

or pharmaceutically acceptable prodrugs, salts, hydrates, solvates, crystal forms or diastereomers thereof, wherein:

$X_1, X_2, X_3, X_4$  are selected from the following:

- (i)  $X_1$  and  $X_2$  are N and  $X_3$  and  $X_4$  are C independently substituted with Y;
- (ii)  $X_1$  and  $X_4$  are N and  $X_2$  and  $X_3$  are C independently substituted with Y;
- (iii)  $X_2$  and  $X_4$  are N and  $X_1$  and  $X_3$  are C independently substituted with Y;
- (iv)  $X_1$  is N and  $X_2, X_3$ , and  $X_4$  are C independently substituted with Y;
- (v)  $X_3$  is N and  $X_1, X_2$ , and  $X_4$  are C independently substituted with Y;
- (vi)  $X_4$  is N and  $X_1, X_2$ , and  $X_3$  are C independently substituted with Y;
- (vii)  $X_2$  is N and  $X_1, X_3$ , and  $X_4$  are C independently substituted with Y; and
- (viii)  $X_1, X_2$  and  $X_3$  are N and  $X_4$  is C substituted with Y;

$R_1$  is H,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkylNR<sub>5</sub>R<sub>6</sub>,  $C_{1-6}$ alkylNR<sub>5</sub>COR<sub>6</sub>,  $C_{1-6}$ alkylNR<sub>5</sub>SO<sub>2</sub>R<sub>6</sub>,  $C_{1-6}$ alkylCO<sub>2</sub>R<sub>5</sub>,  $C_{1-6}$ alkylCONR<sub>5</sub>R<sub>6</sub>, where R<sub>5</sub> and R<sub>6</sub> are each independently H,  $C_{1-4}$ alkyl, aryl, hetaryl,  $C_{1-4}$ alkylaryl,  $C_{1-4}$ alkylhetaryl or may be joined to form an

24.

optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR7;

R7 is selected from H, C<sub>1-4</sub> alkyl;

5 R2 is selected from OH, OC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylOH, OC<sub>2-6</sub>alkylOH, C<sub>1-6</sub>alkylNR8R9, OC<sub>2-6</sub>alkylNR8R9, C<sub>1-6</sub>alkylNR8COR9, OC<sub>2-6</sub>alkylNR8COR9, C<sub>1-6</sub>alkylhetaryl, OC<sub>2-6</sub>alkylhetaryl, OCONR8R9, NR8COOR9, NR10CONR8R9, CONR8R9, NR8COR12;

10 R8, R9 are each independently H, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylNR11R13, hetaryl, cyclohetalkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR14;

R12 is C<sub>2-4</sub>alkyl, C<sub>1-4</sub>alkylNR11R13, hetaryl, cyclohetalkyl;

R11, R13 are each independently H, C<sub>1-4</sub>alkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR14;

15 R14 is selected from H, C<sub>1-4</sub>alkyl;

R10 is H, C<sub>1-4</sub> alkyl;

R3 and R4 are each independently H, halogen, C<sub>1-4</sub>alkyl, OH, OC<sub>1-4</sub>alkyl, CF<sub>3</sub>, OCF<sub>3</sub>;

Q is a bond, or C<sub>1-4</sub>alkyl;

20 W is selected from H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl; where C<sub>1-4</sub>alkyl or C<sub>2-6</sub>alkenyl may be optionally substituted with C<sub>1-4</sub>alkyl, OH, OC<sub>1-4</sub>alkyl, NR15R16;

25 R15, and R16 are each independently H, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyl cycloalkyl, C<sub>1-4</sub>alkyl cyclohetalkyl, aryl, hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR17;

R17 is selected from H, C<sub>1-4</sub>alkyl;

A is aryl, hetaryl optionally substituted with 0-3 substituents independently chosen from halogen, C<sub>1-4</sub> alkyl, CF<sub>3</sub>, aryl, hetaryl, OCF<sub>3</sub>, OC<sub>1-4</sub>alkyl,

25.

OC<sub>2-5</sub>alkylNR<sub>18</sub>R<sub>19</sub>, Oaryl, Ohetaryl, CO<sub>2</sub>R<sub>18</sub>, CONR<sub>18</sub>R<sub>19</sub>, NR<sub>18</sub>R<sub>19</sub>, C<sub>1-4</sub>alkylNR<sub>18</sub>R<sub>19</sub>, NR<sub>20</sub>C<sub>1-4</sub>alkylNR<sub>18</sub>R<sub>19</sub>, NR<sub>18</sub>COR<sub>19</sub>, NR<sub>20</sub>CONR<sub>18</sub>R<sub>19</sub>, NR<sub>18</sub>SO<sub>2</sub>R<sub>19</sub>;

5 R<sub>18</sub>, R<sub>19</sub> are each independently H, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl cyclohetalkyl, aryl, hetaryl, C<sub>1-4</sub>alkyl aryl, C<sub>1-4</sub> alkyl hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>21</sub>;

R<sub>21</sub> is selected from H, C<sub>1-4</sub> alkyl;

R<sub>20</sub> is selected from H, C<sub>1-4</sub> alkyl;

10 Y is selected from H, C<sub>1-4</sub>alkyl, OH, NR<sub>22</sub>R<sub>23</sub>;

R<sub>22</sub>, R<sub>23</sub> are each independently H, C<sub>1-4</sub> alkyl.

In the above description it will be appreciated that:

C<sub>1-4</sub> alkyl means an unsubstituted or optionally substituted straight or branched alkyl chain

15 Aryl means unsubstituted or optionally substituted phenyl or naphthyl.

Hetaryl means an unsubstituted or optionally substituted 5- or 6-membered heteroaromatic ring containing one or more heteroatoms selected from O, N, S.

Cycloalkyl means a 3-8 membered saturated ring

20 Cyclohetalkyl means an optionally substituted 3-8 membered saturated ring containing 1-3 heteroatoms selected from O, S, NR<sub>24</sub>, where R<sub>24</sub> is H, C<sub>1-4</sub> alkyl, aryl, hetaryl.

In a tenth aspect, the present invention provides a composition comprising a carrier and one or more compounds of the ninth aspect.

25 In an eleventh aspect, the present invention provides a method of treatment of a hyperproliferation-related disorder in a subject, the method comprising administering a therapeutically effective amount of at least one of the compounds of the ninth aspect or a composition according to the tenth aspect.

26.

Preferably, the hyperproliferation-related disorder is treatable by the modulation of microtubule polymerisation.

In a twelfth aspect, the present invention consists in a method of treating a protein-kinase related disorder in a subject, the method comprising administering a therapeutically effective amount of at least one of the compounds of the ninth aspect or a composition according to the tenth aspect.

The compounds of this invention include all conformational isomers (eg. cis and trans isomers). The compounds of the present invention have asymmetric centers and therefore exist in different enantiomeric and diastereomeric forms. This invention relates to the use of all optical isomers and stereoisomers of the compounds of the present invention, and mixtures thereof, and to all pharmaceutical compositions and methods of treatment that may employ or contain them. The compounds of formula I, II, III, IV and V may also exist as tautomers. This invention relates to the use of all such tautomers and mixtures thereof.

Compounds of formula I, II, III, IV and V having free amino, amido, hydroxy or carboxylic groups can be converted into prodrugs. Prodrugs include compounds wherein an amino acid residue, or a polypeptide chain of two or more (eg, two, three or four) amino acid residues which are covalently joined through peptide bonds to free amino, hydroxy and carboxylic acid groups of compounds of formulae I, II, III, IV and V. The amino acid residues include the 20 naturally occurring amino acids commonly designated by three letter symbols and also include, 4-hydroxyproline, hydroxylysine, demosine, isodemossine, 3-methylhistidine, norvaline, beta-alanine, gamma-aminobutyric acid, citrulline, homocysteine, homoserine, ornithine and methionine sulfone. Prodrugs also include compounds wherein carbonates, carbamates, amides and alkyl esters which are covalently bonded to the above substituents of formulae I, II, III, IV and V through the carbonyl carbon prodrug sidechain. Prodrugs also include phosphate derivatives of compounds of formulae I, II, III, IV and V (such as acids, salts of acids, or esters) joined through a phosphorus-oxygen bond to a free hydroxyl of compounds of formulae I, II, III, IV and V. Prodrugs also include compounds wherein acyloxyalkyl or phosphonoxyalkyl moieties are covalently attached to compounds of formula I, II, III, IV and V possessing a free hydroxyl group. Acyloxyalkyl or phosphonoxyalkyl moieties may also be covalently attached to compounds of formulae I, II, III, IV and V possessing a pyridyl ring through formation of a N-(acyloxyalkyl)- or N-(phosphonoxyalkyl)-pyridinium salt. This invention also encompasses pharmaceutical compositions containing prodrugs of compounds of the formula I, II, III, IV and V. In addition, this invention encompasses methods of treating or preventing disorders that can be treated or

27.

prevented by the modulation of microtubule dynamics comprising administering prodrugs of compounds of the formula I, II, III, IV and V. This invention further encompasses methods of treating or preventing disorders that can be treated or prevented by the disruption of microtubules comprising administering prodrugs of compounds of the formula II, III, IV and V.

In a still further preferred embodiment the compound possesses *S*chirality at the chiral carbon bearing W, where W is C<sub>1-4</sub> alkyl or C<sub>1-4</sub>alkylamino. The compound can be used as a purified isomer or as a mixture of any ratio of isomers. It is however preferred that the mixture comprises at least 70%, 80%, 90%, 95%, or 99% of the preferred isomer.

10 In a yet further preferred embodiment, the present invention provides a compound selected from the compounds of Table 1; compositions comprising a carrier and one or more compounds of Table 1; and, a method of treatment of a hyperproliferation-related disorder in a subject, the method comprising administering at least one of the compounds of Table 1.

15 More preferably, the hyperproliferation related disorder is treatable by the modulation of microtubule polymerisation.

The present invention further provides for a method of modulating microtubule polymerisation in a subject, the method comprising the administration of a therapeutically effective amount of a compound of any one of Formulas I, II, III, IV and V.

20 In a preferred embodiment, there is provided a method of modulating microtubule polymerisation in a subject, the method comprising the administration of a therapeutically effective amount of a compound selected from the compounds of Table 1.

The present invention provides that the compounds of formula I, II, III, IV and V may be administered for the treatment of hyperproliferation-related disorders. Preferably, the hyperproliferation-related disorder is treatable by the modulation of microtubule polymerisation. In a preferred embodiment, the hyperproliferation-related disorder or disease state is selected from the group consisting of Cancer, such as fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma,



28.

medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms' tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma, and retinoblastoma, and carcinomas forming from tissue of the breast, prostate, kidney, bladder or colon, and neoplastic disorders arising in adipose tissue, such as adipose cell tumors, e.g., lipomas, fibrolipomas, lipoblastomas, lipomatosis, hibernomas, hemangiomas and/or liposarcomas; infectious diseases such as viral, malarial and bacterial infections; vascular restenosis; inflammatory diseases, such as autoimmune diseases, glomerular nephritis myocardial infarction and psoriasis.

The compounds of formula II, II and IV may also be used in the treatment of a hyperproliferation-related disorder or disease state and/or a protein-kinase related disorder or disease state. Preferably, the disease state is selected from the group consisting of Atopy, such as Allergic Asthma, Atopic Dermatitis (Eczema), and Allergic Rhinitis; Cell Mediated Hypersensitivity, such as Allergic Contact Dermatitis and Hypersensitivity Pneumonitis; Rheumatic Diseases, such as Systemic Lupus Erythematosus (SLE), Rheumatoid Arthritis, Juvenile Arthritis, Sjögren's Syndrome, Scleroderma, Polymyositis, Ankylosing Spondylitis, Psoriatic Arthritis; Other autoimmune diseases such as Type I diabetes, autoimmune thyroid disorders, and Alzheimer's disease; Viral Diseases, such as Epstein Barr Virus (EBV), Hepatitis B, Hepatitis C, HIV, HTLV 1, Varicella-Zoster Virus (VZV), Human Papilloma Virus (HPV).

In the treatment or prevention of hyperproliferation-related disorders and/or protein kinase related disorders an appropriate dosage level will generally be about 0.01 to 500 mg per kg patient body weight per day which can be administered in single or multiple doses. Preferably, the dosage level will be about 0.1 to about 250 mg/kg per day; more preferably about 0.5 to about 100 mg/kg per day. A suitable dosage level may be about 0.01 to 250 mg/kg per day, about 0.05 to 100 mg/kg per day, or about 0.1 to 50 mg/kg per day. Within this range the dosage may be 0.05 to 0.5, 0.5 to 5 or 5 to 50 mg/kg per day. For oral administration, the compositions are preferably provided in the form of tablets containing 1.0 to 1000 milligrams of the active ingredient, particularly 1.0, 5.0, 10.0, 15.0, 20.0, 25.0, 50.0, 75.0, 100.0, 150.0, 200.0, 250.0, 300.0, 400.0, 500.0, 600.0, 750.0, 800.0, 900.0, and 1000.0 milligrams of the active ingredient for the symptomatic adjustment of the

29.

dosage to the patient to be treated. The compounds may be administered on a regimen of 1 to 4 times per day, preferably once or twice per day.

It will be understood, however, that the specific dose level and frequency of dosage for any particular patient may be varied and will depend upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of  
5 action of that compound, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular condition, and the host undergoing therapy.

In addition to primates, such as humans, a variety of other mammals can be treated  
10 according to the method of the present invention. For instance, mammals including, but not limited to, cows, sheep, goats, horses, dogs, cats, guinea pigs, rats or other bovine, ovine, equine, canine, feline, rodent or murine species can be treated. However, the method can also be practiced in other species, such as avian species (e.g., chickens).

Diseases and conditions associated with inflammation and infection can be treated using  
15 the method of the present invention. In a preferred embodiment, the disease or condition is one in which the actions of eosinophils and/or lymphocytes are to be inhibited or promoted, in order to modulate the inflammatory response.

The subjects treated in the above methods, in whom which cell growth inhibition is desired, are mammals, including, but not limited to, cows, sheep, goats, horses, dogs, cats,  
20 guinea pigs, rats or other bovine, ovine, equine, canine, feline, rodent or murine species, and preferably a human being, male or female.

The term "therapeutically effective amount" means the amount of the subject composition that will elicit the biological or medical response of a tissue, system, animal or human that is being sought by the researcher, veterinarian, medical doctor or other clinician.

25 The terms "administration of" and or "administering a" compound should be understood to mean providing a compound of the invention to the individual in need of treatment.

The present invention provides pharmaceutical compositions comprising at least one of the compounds of the formula I, II, III, IV and V capable of treating a hyperproliferation-related disorder in an amount effective therefore, and a pharmaceutically acceptable  
30 vehicle or diluent. The compositions of the present invention may contain other therapeutic agents as described below, and may be formulated, for example, by employing conventional solid or liquid vehicles or diluents, as well as pharmaceutical

30.

additives of a type appropriate to the mode of desired administration (for example, excipients, binders, preservatives, stabilizers, flavors, etc.) according to techniques such as those well known in the art of pharmaceutical formulation.

5 The compounds of the formula I, II, III, IV and V may be administered by any suitable means, for example, orally, such as in the form of tablets, capsules, granules or powders; sublingually; buccally; parenterally, such as by subcutaneous, intravenous, intramuscular, or intracisternal injection or infusion techniques (e.g., as sterile injectable aqueous or non-aqueous solutions or suspensions); nasally such as by inhalation spray; topically, such as in the form of a cream or ointment; or rectally such as in the form of suppositories; in  
10 dosage unit formulations containing non-toxic, pharmaceutically acceptable vehicles or diluents. The compounds may, for example, be administered in a form suitable for immediate release or extended release. Immediate release or extended release may be achieved by the use of suitable pharmaceutical compositions comprising the present compounds, or, particularly in the case of extended release, by the use of devices such as  
15 subcutaneous implants or osmotic pumps.

The term "composition" as used herein is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts. By "pharmaceutically acceptable" it is meant the carrier, diluent or excipient  
20 must be compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

The pharmaceutical compositions for the administration of the compounds of this invention may conveniently be presented in dosage unit form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of  
25 bringing the active ingredient into association with the carrier which constitutes one or more accessory ingredients. In general, the pharmaceutical compositions are prepared by uniformly and intimately bringing the active ingredient into association with a liquid carrier or a finely divided solid carrier or both, and then, if necessary, shaping the product into the desired formulation. In the pharmaceutical composition the active object  
30 compound is included in an amount sufficient to produce the desired effect upon the process or condition of diseases. As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

31.

The pharmaceutical compositions containing the active ingredient may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. They may also be coated to form osmotic therapeutic tablets for control release.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more



32.

preservatives, for example ethyl, or n-propyl, p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

5 Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

10 Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be  
15 present.

The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally- occurring gums, for example gum acacia or gum tragacanth, naturally-  
20 occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavoring agents.

25 Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents.

The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension. This suspension may be formulated according to the known art  
30 using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butane diol. Among the acceptable vehicles and solvents that

33.

may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the  
5 preparation of injectables.

The compounds of the present invention may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to  
10 release the drug. Such materials are cocoa butter and polyethylene glycols.

For topical use, creams, ointments, jellies, solutions or suspensions, etc., containing the compounds of the present invention are employed. (For purposes of this application, topical application shall include mouthwashes and gargles.)

The compounds of the present invention can also be administered in the form of  
15 liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multilamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically acceptable and metabolisable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to a compound of the  
20 present invention, stabilisers, preservatives, excipients and the like. The preferred lipids are the phospholipids and phosphatidyl cholines, both natural and synthetic. Methods to form liposomes are known in the art.

The pharmaceutical composition and method of the present invention may further comprise other therapeutically active compounds as noted herein which are usually  
25 applied in the treatment of the above mentioned pathological conditions. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve  
30 therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

Examples of other therapeutic agents include the following:



34.

Tyrosine kinase inhibitors, such as imatinib (Glivec™), and gefitinib (Iressa™) *inter alia*; cyclosporins (e.g., cyclosporin A), CTLA4-Ig, antibodies such as ICAM-3, anti-IL-2 receptor (Anti-Tac), anti-CD45RB, anti-CD2, anti-CD3 (OKT-3), anti-CD4, anti-CD80, anti-CD86, agents blocking the interaction between CD40 and gp39, such as antibodies  
 5 specific for CD40 and/or gp39 (i.e., CD154), fusion proteins constructed from CD40 and gp39 (CD401g and CD8gp39), inhibitors, such as nuclear translocation inhibitors, of NF-kappa B function, such as deoxyspergualin (DSG), cholesterol biosynthesis inhibitors such as HMG CoA reductase inhibitors (lovastatin and simvastatin), non-steroidal antiinflammatory drugs (NSAIDs) such as ibuprofen, aspirin, acetaminophen and  
 10 cyclooxygenase inhibitors such as rofecoxib, steroids such as prednisolone or dexamethasone, gold compounds, antiproliferative agents such as methotrexate, FK506 (tacrolimus, Prograf), mycophenolate mofetil, antineoplastic agents such as azathioprine, VP-16, etoposide, fludarabine, cisplatin, doxorubicin, adriamycin, amsacrine, camptothecin, cytarabine, gemcitabine, fluorodeoxyuridine, melphalan and  
 15 cyclophosphamide, TNF- $\alpha$  inhibitors such as tenidap, anti-TNF antibodies or soluble TNF receptor, and rapamycin (sirolimus or Rapamune) or derivatives thereof.

When other therapeutic agents are employed in combination with the compounds of the present invention they may be used for example in amounts as noted in the Physician Desk Reference (PDR) or as otherwise determined by one of ordinary skill in the art.

20 The pharmaceutical composition and method of the present invention may further comprise other therapeutically active compounds as noted herein which are known inhibitors or substrates of drug efflux systems or drug detoxification and excretory systems. Such systems include P-glycoprotein, multidrug resistance-associated protein, lung resistance protein and glutathione S-transferase isoenzymes alpha, mu, pi, sigma,  
 25 theta, zeta and kappa. Co-administration of drugs known to inhibit or reduce the activity of these systems may increase the efficacy of the compounds described in the present invention through increasing the amount of therapeutic agent in the cell. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages, thus reducing the potential for adverse side effects. Examples of inhibitors or substrates for  
 30 these systems include; verapamil, probenecid, dipyridamole, ethacrynic acid, indomethacin, sulfasalazine, buthionine sulfoximine, cyclosporin A and tamoxifen.

Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or

35.

step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

All publications mentioned in this specification are herein incorporated by reference.

5 Any discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is solely for the purpose of providing a context for the present invention. It is not to be taken as an admission that any or all of these matters form part of the prior art base or were common general knowledge in the field relevant to the present invention as it existed in Australia before the priority date of each claim of this application.

10 In order that the nature of the present invention may be more clearly understood preferred forms thereof will now be described by reference to the following non-limiting Examples.

36.

**EXAMPLES****MATERIALS AND METHODS:***Compound Synthesis*

Compounds are generally prepared in a 2-step process starting from a dihaloheterocycle.

- 5 The first step is a nucleophilic aromatic substitution to generate a monoamino-monohalo intermediate.

The nucleophilic aromatic substitution is typically carried out by addition of a primary or secondary amine to the di-halogenated heterocycle in a solvent such as ethanol, isopropanol, tert-butanol, dioxane, THF, DMF, ethoxyethanol, toluene or xylene. The  
10 reaction is typically performed at elevated temperature in the presence of excess amine or a non-nucleophilic base such as triethylamine or diisopropylethylamine, or an inorganic base such as potassium carbonate or sodium carbonate.

Alternatively, the amino substituent may be introduced through a transition metal catalysed amination reaction. Typical catalysts for such transformations include  
15  $\text{Pd}(\text{OAc})_2/\text{P}(\text{t-Bu})_3$ ,  $\text{Pd}_2(\text{dba})_3/\text{BINAP}$  and  $\text{Pd}(\text{OAc})_2/\text{BINAP}$ . These reactions are typically carried out in solvents such as toluene or dioxane, in the presence of bases such as caesium carbonate or sodium or potassium tert-butoxide at temperatures ranging from room temperature to reflux.

The amines employed in the first step of the synthesis of these compounds are obtained  
20 commercially or are prepared using methods well known to those skilled in the art. Of particular interest are  $\alpha$ -alkylbenzylamines which may be prepared through reduction of oximes. Typical reductants include lithium aluminium hydride, hydrogen gas in the presence of palladium on charcoal catalyst, Zn in the presence of hydrochloric acid, sodium borohydride in the presence of a Lewis acid such as  $\text{TiCl}_3$ ,  $\text{ZrCl}_4$ ,  $\text{NiCl}_2$  and  $\text{MoO}_3$ ,  
25 or sodium borohydride in conjunction with Amberlyst H15 ion exchange resin and LiCl.

$\alpha$ -Alkylbenzylamines may also be prepared by reductive amination of the corresponding ketones. A classical method for such a transformation is the Leuckart-Wallach reaction, though catalytic conditions ( $\text{HCO}_2\text{NH}_4$ ,  $[(\text{CH}_3)_5\text{C}_5\text{RhCl}_2]_2$ ) or alternative procedures (e.g.  $\text{NH}_4\text{OAc}$ ,  $\text{Na}(\text{CN})\text{BH}_3$ ) can also be used.

37.

$\alpha$ -Alkylbenzylamines may also be prepared from the corresponding  $\alpha$ -alkylbenzyl alcohols. Such methods include derivatisation of the hydroxyl as a mesylate or tosylate and displacement with a nitrogen nucleophile, such as phthalimide or azide which is converted to the primary amine using conventional synthetic methods; or, displacement  
5 of the hydroxyl with a suitable nitrogen nucleophile under Mitsunobu-like conditions.  $\alpha$ -Alkylbenzyl alcohols can be prepared by reduction of the corresponding ketones with a reducing agent such as sodium borohydride in a solvent such as methanol. Alternatively,  $\alpha$ -alkylbenzyl alcohols can be obtained through addition of an alkyl metal species (such as a Grignard reagent) to a benzaldehyde derivative, typically performed at room  
10 temperature or below in solvents such as tetrahydrofuran.

$\alpha$ -Alkyl benzylamines of high optical purity may be prepared from chiral  $\alpha$ -alkyl benzyl alcohols using the methods outlined above. The chiral  $\alpha$ -alkyl benzyl alcohols may be obtained through chiral reduction of the corresponding ketones. Chiral reducing methods are now well known in organic chemistry and include enzymatic processes,  
15 asymmetric hydrogenation procedures and chiral oxazaborolidines.

The second step of the synthesis typically involves a palladium mediated cross-coupling of the monoamino-monochloro intermediate with a suitably functionalised coupling partner. Typical coupling partners are boronic acids or esters (Suzuki coupling: see for example Miyaura, N. and Suzuki, *Chem Rev.* **1995**, 952457), stannanes (Stille coupling:  
20 see for example Stille, J.K., *Angew. Chem., Int. Ed. Engl.*, **1986**, 25, 508), Grignard reagents (Kumada coupling: Kumada, M.; Tamao, K.; Sumitani, K. *Org. Synth.* **1988**, Coll. Vol.6, 407.) or organozinc species (Negishi coupling: Negishi, E.; *J. Organomet. Chem.* **2002**, 653, 34).

The Suzuki coupling is the preferred coupling method and is typically performed in a  
25 solvent such as DME, THF, DMF, ethanol, propanol, toluene, or 1,4-dioxane in the presence of a base such as potassium carbonate, lithium hydroxide, caesium carbonate, sodium hydroxide, potassium fluoride or potassium phosphate. The reaction may be carried out at elevated temperatures and the palladium catalyst employed may be selected from  $\text{Pd}(\text{PPh}_3)_4$ ,  $\text{Pd}(\text{OAc})_2$ ,  $[\text{PdCl}_2(\text{dppf})]$ ,  $\text{Pd}_2(\text{dba})_3/\text{P}(\text{t-Bu})_3$ .

30 The products formed from this reaction sequence may be further derivatised using techniques well-known to those skilled in the art. Alternatively, derivatisation of the mono-amino mono-halo intermediate may be undertaken prior to displacement of the halo substituent. This derivatisation typically involves functionality originally present on the amine species and employs methods well known to those skilled in the art.

38.


Alternatively the preparation sequence can be reversed beginning with the palladium mediated cross-coupling reaction to furnish a mono-halo heterocyclic species. Amine displacement of the halo substituent can then be undertaken using procedures outlined above.

- 5 In order that the nature of the present invention may be more clearly understood, preferred forms thereof will now be described with reference to the following non-limiting examples.

### EXAMPLE 1

#### 1-[4-(Trifluoromethyl)phenyl]butan-1-ol

10



15


A 2M solution of propylmagnesium chloride in ether (4 ml, 8 mmol) was added to a solution of the aldehyde (1.14 g, 6.6 mmol) in dry THF (10 ml) cooled to 0 °C under N<sub>2</sub>. The mixture was stirred for 16h at room temperature, after which time saturated ammonium chloride solution was added. The product was extracted into ethyl acetate, and the ethyl acetate layer dried and concentrated to furnish pure product (1.4g, 98%).

<sup>1</sup>H-n.m.r. (CDCl<sub>3</sub>) δ 0.94 (t, 3H, *J* = 7.2Hz, CH<sub>3</sub>), 1.41 (m, 2H, CH<sub>2</sub>), 1.75 (m, CH<sub>2</sub>, 2H), 4.77 (br s, 1H, CH), 7.44-7.62(m, 4H, ArH)

### EXAMPLE 2

#### 1-(1-Azidobutyl)-4-(trifluoromethyl)benzene

20



A solution of 1-[4-(trifluoromethyl)phenyl]butan-1-ol (1.4 g, 6.4 mmol) and diphenylphosphoryl azide (2.8 ml, 12.8 mmol) in THF (6 mL) cooled to -10 °C under N<sub>2</sub> was treated with DBU (1.9 mL, 12.8 mmol). The resulting solution was stirred at room temperature for 20 hours and then diluted with a mixture of ether and H<sub>2</sub>O. The organic




39.

phase was dried and concentrated and the residue purified by column chromatography using hexane:ethyl acetate (10:1) as eluent to furnish pure azide (0.85 g, 54%).

<sup>1</sup>H-n.m.r. (CDCl<sub>3</sub>) δ 0.94 (t, 3H, *J* = 7.2 Hz, CH<sub>3</sub>), 1.37 (m, 2H, CH<sub>2</sub>), 1.75 (m, 2H, CH<sub>2</sub>), 4.50 (t, 1H, CH), 7.42 (d, *J* = 7.8 Hz, 2H, ArH), 7.64 (d, 2H, *J* = 7.8 Hz, ArH)

### 5 EXAMPLE 3

#### *1-[4-(Trifluoromethyl)phenyl]butan-1-amine*




A mixture of 1-(1-azidobutyl)-4-(trifluoromethyl)benzene (0.84 g, 3.5 mmol) and triphenylphosphine (1.8g, 6.9 mmol) in ethyl acetate (6 mL) and 10% HCl (6ml) was stirred at room temperature for 64 h. The aqueous phase was collected and the organic phase extracted with 10% HCl (3 x 5mL). The aqueous layers were combined and basified with 5M NaOH, and then extracted with ethyl acetate (5 x 15mL). The organic phase was dried and concentrated to give pure amine (0.4 g, 54%).

<sup>1</sup>H-n.m.r. (CDCl<sub>3</sub>) δ 0.91 (t, *J* = 7.4 Hz, 3H CH<sub>3</sub>), 1.31 (m, 2H, CH<sub>2</sub>), 1.62 (m, 2H, CH<sub>2</sub>), 3.97 (m, 1H, CH), 7.43 (AA'XX', 2H, ArH), 7.58 (AA'XX', 2H, ArH)

### EXAMPLE 4

#### *3-Fluoro-N-methoxy-N-methylbenzamide*



To a suspension of 3-fluorobenzoic acid (140 mg, 1 mmol) and *N,N*-dimethylhydroxylamine hydrochloride (107 mg, 1.1 mmol) in dichloromethane (2.5 mL) was added 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (EDC) (211 mg, 1.1 mmol) and the mixture stirred at room temperature for 75 h. The solvents were removed under reduced pressure and the residue chromatographed using ethyl acetate-hexane (4:6) to separate the pure product (130 mg, 71%).




40.

$^1\text{H}$ -n.m.r. ( $\text{CDCl}_3$ )  $\delta$  3.36 (s, 3H, N-Me), 3.55 (s, 3H, N-OMe), 7.1.-7.2 (m, 1H, ArH), 7.3-7.5 (m, 3H, Ar)


**EXAMPLE 5***1-(3-Fluorophenyl)butan-1-one*

5



To a solution of 3-fluoro-*N*-methoxy-*N*-methylbenzamide (130 mg, 0.71 mmol) in dry THF (2 mL) cooled to  $-10^\circ\text{C}$  was added propyl magnesium chloride (532  $\mu\text{l}$ , 2M solution in ether, 1.1 mmol) under nitrogen. The solution was stirred at  $-10^\circ\text{C}$  for 1h and at room temperature for 75 min. The solution was the poured into saturated aqueous ammonium chloride and the product extracted into ethyl acetate (3 x 25 mL). The organic layers were combined, washed with brine, dried ( $\text{MgSO}_4$ ) and concentrated to give a pale yellow oil, which was purified by chromatography using ethyl acetate-hexane (1:9) to separate the pure product (78 mg, 66%).

$^1\text{H}$ -n.m.r. ( $\text{CDCl}_3$ )  $\delta$  1.01 (t, 3H,  $J=7.4$  Hz, Me), 1.77 (m, 2H,  $\text{CH}_2\text{CH}_3$ ), 2.93 (t,  $J=7.2$  Hz, 2H,  $\text{COCH}_2$ ), 7.15-7.30 (m, 1H, Ar), 7.35-7.50 (m, 1H, Ar), 7.60-7.70 (m, 1H, Ar), 7.70-7.80 (m, 1H, Ar).

**EXAMPLE 6***1-(5-Methylpyridin-3-yl)butan-1-amine*

To a solution of 1-(5-methylpyridin-3-yl)butan-1-one (800 mg, 4.9 mmol) and ammonium formate (1.55 g, 24.5 mmol) in methanol (5 mL) under nitrogen was added dichloro(pentamethylcyclopentadienyl)rhodium (III) dimer (45 mg, 0.074 mmol). The solution was heated at reflux for 8 h. after which time the solution was cooled to room temperature and acidified to pH ~2 with 2M HCl. The mixture was washed with dichloromethane (3 x 15 mL) and the aqueous phase then basified to pH ~12 by addition

41.


of solid KOH. The aqueous phase was extracted with dichloromethane (3 x 15 mL) and the combined organic layers washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to give a pure product (620 mg, 77%).

<sup>1</sup>H-n.m.r. (CDCl<sub>3</sub>) δ 0.91 (t, 3H, *J* = 7.4 Hz, CH<sub>2</sub> CH<sub>2</sub> CH<sub>3</sub>), 1.14-1.82 (m, 6H, 2 x CH<sub>2</sub>, NH),  
 5 2.33 (s, 3H, CH<sub>3</sub>), 3.91 (t, 1H, *J* = 6.8 Hz, CHNH<sub>2</sub>), 7.47 (s, 1H, ArH), 8.32 (br s, 2H, Ar).

**EXAMPLE 7***(1R)-1-(4-Fluorophenyl)butan-1-ol*

Diethylaniline-borane complex (2.4 mL, 13.4 mmol) was added dropwise over 5 min. to a  
 10 solution of (1S,2R)-1-aminoindan-2-ol (201 mg, 1.4 mmol) in dry THF (20 mL) cooled to 0°  
 C under nitrogen. After 40 min. a solution of the ketone in THF (40 mL) was added  
 dropwise over 90 min. The cooling bath was then removed and the solution then stirred  
 at RT for 5h. Acetone (6 mL) was added and the resulting solution stirred at RT for 60  
 15 min. before the solvents were removed under reduced pressure. The residue was  
 dissolved in toluene (100 mL) and washed successively with 1M H<sub>2</sub>SO<sub>4</sub> (4 x 50 mL), H<sub>2</sub>O  
 (2 x 50 mL) and brine (50 mL) and then dried (Na<sub>2</sub>SO<sub>4</sub>).

<sup>1</sup>H-n.m.r. (CDCl<sub>3</sub>) δ 0.93 (t, 3H, *J* = 7.2 Hz, CH<sub>2</sub> CH<sub>2</sub> CH<sub>3</sub>), 1.17-1.51 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>),  
 1.58-1.82 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 4.66 (t, *J* = 6.6 Hz, CHNH<sub>2</sub>), 6.98-7.07 (m, 2H, ArH), 7.26-  
 7.34 (m, 2H, ArH).

**EXAMPLE 8***(1R)-1-Pyridin-3-ylbutan-1-ol*

To a solution of S-(-)- α, α -diphenyl-2-pyrrolidinemethanol (500 mg, 2 mmol) and  
 trimethylborate (0.27ml, 2.4 mmol) in THF (20 mL) was added borane-dimethylsulfide  
 25 complex (20mL, 2M in THF, 40 mmol). The mixture was cooled to 0°C and 1-pyridin-3-


42.

ylbutan-1-one was added dropwise over 4 h. The solution was stirred at RT overnight and then treated with 2M HCl (175 mL) and stirring continued for 4 h. The volatiles were removed under reduced pressure and the resulting cloudy solution basified to pH 11 with aqueous ammonia. The product was extracted into ethyl acetate (3 x 100 mL) and the combined organic layers washed with brine (100 mL) and dried (Na<sub>2</sub>SO<sub>4</sub>). Removal of the solvent under reduced pressure afforded ~3g of product.

<sup>1</sup>H-n.m.r. (CDCl<sub>3</sub>) δ 0.94 (t, 3H, *J*=7.4 Hz, CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 1.20-1.54 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 1.60-1.91 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 2.64 (br s, 1H, OH), 4.73 (t, 1H, *J*=6.6 Hz, CHOH), 7.24-7.30 (m, 1H, Ar), 7.68-7.74 (m, 1H, Ar), 8.47-8.52 (m, 2H, Ar).

## 10 EXAMPLE 9

### *3-Amino-3-phenylpropan-1-ol*




To a suspension of 3-amino-3-phenylpropanoic acid (2.0 g, 12.1 mmol) in dry THF (45 mL) cooled to 0 °C under N<sub>2</sub> was added portionwise over 20 minutes solid LiAlH<sub>4</sub> (920 mg, 24.2 mmol). Stirring was continued at room temperature for 24 h. after which time solid Na<sub>2</sub>SO<sub>4</sub>·10H<sub>2</sub>O was added with stirring until only a heavy white precipitate was present. The organic layer was diluted with ether and filtered through Celite<sup>®</sup> and the concentrated *in vacuo*. The residue was dissolved in ethyl acetate (50 mL) and extracted with 1N HCl (3 x 40 mL). The aqueous layers were combined and basified to pH~12 with 5M NaOH. The aqueous phase was extracted with ethyl acetate (3 x 50 mL) and the combined organic phases dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated *in vacuo* to separate the product which was used without further purification (0.9g, 49%).

<sup>1</sup>H-n.m.r. (CDCl<sub>3</sub>) δ 1.84-1.94 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>OH), 2.68 (br s, 1H, NH<sub>2</sub>), 3.79 (t, 2H, *J*= 5.8 Hz, CH<sub>2</sub>CH<sub>2</sub>OH), 4.08-4.15 (m, 1H, CHNH<sub>2</sub>), 4.77 (q, 1H, CH), 7.21-7.38 (m, 5H, ArH)

## EXMAPLE 10

### *6-Chloro-N-[1-(4-methylphenyl)butyl]pyrazin-2-amine*

43.




To a solution of 1-[4-(trifluoromethyl)phenyl]butan-1-amine (0.40 g, 1.9 mmol) and 2,6-dichloropyrazine (0.55 g, 3.7 mmol) in 1,4-dioxane (6 mL) was added anhydrous potassium carbonate (0.39 g, 2.8 mmol). The resulting mixture was heated at reflux for 18 h. After cooling to room temperature, the mixture was diluted with ethyl acetate and H<sub>2</sub>O. The organic phase was collected, dried and concentrated. The residue was purified by flash chromatography eluting with ethyl acetate-hexane (1:1) to give pure product (0.03 g, 5%).

<sup>1</sup>H-n.m.r. (CDCl<sub>3</sub>) δ 0.95 (t, 3H, *J* = 7.4 Hz, CH<sub>3</sub>), 1.39 (m, 2H, CH<sub>2</sub>), 1.81 (m, 2H, CH<sub>2</sub>), 4.77 (q, 1H, CH), 5.09 (br d, 1H, NH), 7.42–7.62 (m, 5H, ArH), 7.80 (s, 1H, pyraz.-H)

#### EXAMPLE 11


*N*-(3-[1-[(6-Chloropyrazin-2-yl)amino]butyl]phenyl)acetamide



To a stirred solution of *N*-[1-(3-aminophenyl)butyl]-6-chloropyrazin-2-amine (0.10 g, 0.36 mmol) and triethylamine (100 μL, 0.72 mmol) in dichloromethane (3 mL) cooled to 0 °C was added acetyl chloride (31 μL, 0.43 mmol). The resulting mixture was stirred at room temperature for 18 h. then diluted with dichloromethane (10 mL) and washed with H<sub>2</sub>O (10 mL) and brine (10 mL). The organic phase was collected, dried and concentrated and the residue purified by flash chromatography eluting with ethyl acetate-hexane (1:1) to give pure product (94 mg, 82%).

<sup>1</sup>H-n.m.r. (CDCl<sub>3</sub>) δ 0.93 (t, 3H, *J* = 7.4 Hz, Me), 1.28–1.43 (m, 2H, CH<sub>2</sub>), 1.76–1.85 (m, 2H, CH<sub>2</sub>), 2.16 (s, 3H, COCH<sub>3</sub>), 4.48–4.59 (m, 1H, CHCH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 5.15 (br d, *J* = 6.8 Hz, 1H, NH), 7.06 (d, 1H, *J* = 7.4 Hz, Ar), 7.23–7.33 (m, 3H, Ar), 7.54 (s, 1H, CONH), 7.60 (s, 1H, pyraz.-H), 7.77 (s, 1H, pyraz.-H).


44.

**EXAMPLE 12***N*-[2-Methoxy-4-(6-[(1*S*)-1-pyridin-3-ylbutyl]amino)pyrazin-2-yl]phenyl]-*N'*-ethylurea

5 Under a nitrogen atmosphere a mixture of 6-chloro-*N*-[(1*S*)-1-pyridin-3-ylbutyl]pyrazin-2-amine (53 g, 0.2 mmol), 4-[(ethylamino)carbonylamino]-3-methoxyphenylboronic acid pinacol diester (69 mg, 0.23 mmol), tetrakis(triphenylphosphine)palladium(0) (23 mg, 0.02 mmol) in toluene—*n*-propanol (2.6 mL, 3:1) was treated with 2M aqueous sodium carbonate solution (0.15 mL, 0.3 mmol). The resulting mixture was stirred vigorously

10 whilst being heated at 100°C for 22 hours. Once cool ethyl acetate (10 mL) was added and the mixture washed with H<sub>2</sub>O (6 x 10 mL) and brine (10 mL) and then dried (Na<sub>2</sub>SO<sub>4</sub>). Removal of solvent *in vacuo* then yielded crude product, which was purified by column chromatography using dichloromethane-methanol-aqueous ammonia (93:7:1) as eluent to furnish the product (60 mg).

15 <sup>1</sup>H-n.m.r. (CDCl<sub>3</sub>) δ 0.98 (t, 3H, *J* = 7.4 Hz, CH<sub>3</sub>), 1.14 (t, 3H, *J* = 7.2 Hz, CH<sub>3</sub>), 1.36-1.52 (m, 2H, CH<sub>2</sub>), 1.78-1.98 (m, 2H, CH<sub>2</sub>), 2.27 (s, 3H, CH<sub>3</sub>), 3.23-3.37 (m, 2H, CH<sub>2</sub>), 4.84-4.94 (m, 2H, CH, NHCONH), 5.19 (d, 1H, *J* = 6.2 Hz, CHNH), 6.31 (s, 1H, ArNH), 7.24-7.30 (m, 1H, ArH), 7.55-7.73 (m, 4H, ArH), 7.75 (s, 1H, pyraz.-H), 8.21 (s, 1H, pyraz.-H), 8.49-8.52 (m, 1H, ArH), 8.69 (d, 1H, *J* = 2.0 Hz, ArH).

**EXAMPLE 13***4-[(Ethylamino)carbonylamino]-3-methylphenylboronic acid pinacol diester*

45.


The aniline (500mg, 2.7 mmol) was dissolved in pyridine (5 mL) and to this was added ethyl isocyanate (424  $\mu$ L, 5.4 mmol). The resulting solution was stirred overnight during which time a thick precipitate formed which was filtered and dried *in vacuo* to give the ethyl urea (595 mg, 86%).  $^1\text{H}$ -n.m.r. ( $\text{d}_6$ -DMSO)  $\delta$  1.05 (t, 3H,  $J$ = 7.4 Hz,  $\text{CH}_3$ ), 2.15 (s, 3H,  $\text{CH}_3$ ), 3.09 (dq, 2H,  $J$ = 7.4, 5.4 Hz,  $\text{CH}_2$ ), 6.55 (br t, 1H,  $J$ = 5.4 Hz, NH), 7.24 (dd, 1H,  $J$ = 8.8, 2.2 Hz, Ar-H), 7.30 (d, 1H,  $J$ = 2.2 Hz, Ar-H), 7.66 (s, 1H, NH), 7.82 (d, 1H,  $J$ = 8.8 Hz, ArH).

To a degassed suspension of the ethyl urea bromide (514 mg, 2 mmol), potassium acetate (784 mg, 8 mmol) and bis(pinacolato)diboron (508 mg, 2 mmol) in a mixture of ethanol (10 mL) and dioxane (2 mL) was added dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (II) dichloromethane adduct (44 mg, 0.06 mmol). The solution was heated at reflux for 2 h. and then allowed to cool to RT. The solvent was removed under reduced pressure and the residue dissolved in dichloromethane (60 mL). This was washed with  $\text{H}_2\text{O}$  (2 x 30 mL) and brine (30 mL), dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated in vacuo. The residue was chromatographed using ethyl acetate-hexane (1:1) as eluant to separate the pure product as a solid (351 mg, 58%).

$^1\text{H}$ -n.m.r. ( $\text{CDCl}_3$ )  $\delta$  1.12 (t, 3H,  $J$ = 6.8 Hz,  $\text{CH}_3$ ), 1.34 (s, 12H,  $\text{CH}_3$ ), 2.26 (s, 3H,  $\text{CH}_3$ ), 3.09 (dq, 2H,  $J$ = 6.8, 5.2 Hz,  $\text{CH}_2$ ), 4.69 (br t, 1H, NH), 6.06 (s, 1H, NH), 7.49 (d, 1H,  $J$ = 7.8 Hz, Ar-H), 7.67 (m, 2H, ArH).

## 20 EXAMPLE 14

### *5-Bromo-N-(1-phenylbutyl)pyridin-3-amine*




To a degassed solution of 3,5-dibromopyridine (0.4 g, 1.7 mmol), sodium tert-butoxide (227 mg, 3.6 mmol), tris(dibenzylideneacetone)dipalladium (145 mg, 0.09 mmol) and *rac*-BINAP (105 mg, 0.2 mmol) in dry toluene (10 mL) was added 1-phenylbutan-1-amine (0.2 mL). The mixture was heated at reflux for 24h and upon cooling to room temperature was diluted with ether and washed with brine (3 x 30 mL). The solution was dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated and the residue chromatographed using ethyl acetate-hexane (40:60) as eluant. From the early fractions 97 mg of pure product was obtained.




46.

<sup>1</sup>H-n.m.r. (CDCl<sub>3</sub>) δ 0.94 (t, 3H, *J* = 7.2 Hz, CH<sub>3</sub>), 1.21-1.53 (m, 2H, CH<sub>2</sub>), 1.68-1.87 (m, 2H, CH<sub>2</sub>), 4.19-4.32 (m, 2H, CH and NH), 6.86 (t, 1H, *J* = 2.0 Hz, ArH), 7.20-7.38 (m, 5H, ArH), 7.88 (d, 1H, *J* = 2.0 Hz, ArH), 7.92 (d, 1H, *J* = 2.0 Hz, ArH).

**EXAMPLE 15**5 *5-[4-(Benzyloxy)-3-methoxyphenyl]-3-chloro-6-methyl-1,2,4-triazine*

A degassed solution of 3,5-dichloro-6-methyl-1,2,4-triazine (151 mg, 0.9 mmol), 4-(benzyloxy)-3-methoxybenzene boronic acid (236 g, 0.9 mmol), tetrakis(triphenylphosphine)palladium(0) (55 mg, 0.05 mmol) and tripotassium phosphate  
 10 (249 mg, 1.2 mmol) in dimethoxyethane (5 mL) was heated at reflux for 24 h. Upon cooling to room temperature the solution was diluted with CHCl<sub>3</sub> and filtered through Celite<sup>®</sup>. The filtrate was concentrated *in vacuo* and the product purified by column chromatography using ethyl acetate-hexane (25:75) as eluent to furnish (75 mg, 22%).

<sup>1</sup>H-n.m.r. (CDCl<sub>3</sub>) δ 0.88 (s, 3H, Ar-CH<sub>3</sub>), 3.98 (s, 3H, OCH<sub>3</sub>), 5.26 (s, 2H, CH<sub>2</sub>), 6.99 (d, 1H, *J* = 8.5 Hz, ArH), 7.25-7.44 (m, 7H, ArH).  
 15

**EXAMPLE 16***5-[4-(Benzyloxy)-3-methoxyphenyl]-6-methyl-N-(1-phenylpropyl)-1,2,4-triazin-3-amine*

A solution of the triazene (50 mg, 0.14 mmol) in ethoxyethanol (2.5 mL) was treated with diisopropylamine (50 μL, 0.28 mmol) and 1-phenylbutan-1-amine (46 μL, 0.28 mmol) and the resulting solution heated at 110°C for 3 days. The solvent was then removed *in vacuo* to give an orange residue which was chromatographed using ethyl acetate-  
 20


47.

dichloromethane (20:80) as eluant to separate the product as a pale yellow solid (60 mg, 90%).

<sup>1</sup>H-n.m.r. (CDCl<sub>3</sub>) δ 0.94 (t, 3H, CH<sub>3</sub>), 1.21-1.53 (m, 2H, CH<sub>2</sub>), 1.78-1.97 (m, 2H, CH<sub>2</sub>), □.62 (s, 3H, Ar-CH<sub>3</sub>), 3.89 (s, 3H, OCH<sub>3</sub>), 5.10 (br q, 1H, CH<sub>2</sub>), 5.22 (s, 2H, CH<sub>2</sub>), 5.65 (br s, 1H, NH), 6.87 (d, 1H, *J* = 9.2 Hz, ArH), 7.12-7.44 (m, 12H, Ar-H).

**EXAMPLE 17**

*2-Chloro-5-methyl-N-[(1S)-1-pyridin-3-ylbutyl]pyrimidin-4-amine*




To a solution of (1S)-1-pyridin-3-ylbutan-1-amine (600 mg, 4.0 mmol) in ethanol (20 mL) was added 2,4-dichloro-5-methylpyrimidine (717 mg, 4.4 mmol) and diisopropylethylamine (1.4 mL, 8mmol). The solution was heated at reflux for 24h after which time the solvent was removed under reduced pressure. The residue was dissolved in ethyl acetate (30 mL) and washed with H<sub>2</sub>O (2 x 15 mL), brine (15 mL) and dried (Na<sub>2</sub>SO<sub>4</sub>). The residue remaining after concentration *in vacuo* was chromatographed using ethyl acetate-hexanes (9:1) as eluant to separate the desired product (370 mg).

<sup>1</sup>H-n.m.r. (CDCl<sub>3</sub>) δ 0.96 (t, 3H, *J* = 7.2 Hz, CH<sub>3</sub>), 1.23-1.52 (m, 2H, CH<sub>2</sub>), 1.85-1.99 (m, 2H, CH<sub>2</sub>), 2.03 (d, 3H, *J* = 0.8 Hz, Ar-Me), 5.03 (br d, 1H, *J* = 7.2 Hz, NH), 5.22-5.33 (m, 1H, CH), 7.24-7.30 (m, 1H, ArH), 7.65-7.71 (m, 1H, ArH), 7.81 (d, 1H, *J* = 0.8 Hz, ArH), 8.51 (dd, *J* = 5.6, 1.4 Hz, 1H, ArH), 8.64 (d, 1H, *J* = 1.8 Hz, ArH).

**EXAMPLE 18**

*N-Ethyl-N'-[2-methoxy-4-(5-methyl-4-[(1S)-1-pyridin-3-ylbutyl]amino)pyrimidin-2-yl]phenyl]urea*

48.




Under a nitrogen atmosphere a mixture of 2-chloro-5-methyl-N-[(1S)-1-pyridin-3-ylbutyl]pyrimidin-4-amine (277 mg, 1.0 mmol), 4-[[[(ethylamino)carbonyl]amino]-3-methoxyphenyl]boronic acid pinacol diester (416 mg, 1.3 mmol),  
 5 tetrakis(triphenylphosphine)palladium(0) (116 mg, 0.1 mmol) in toluene—n-propanol (12 mL, 3:1) was treated with 2M aqueous sodium carbonate solution (750  $\mu$ L, 1.5 mmol). The resulting mixture was stirred vigorously whilst being heated at 100 °C for 17 hours. Once cool ethyl acetate (25 mL) was added and the mixture washed with H<sub>2</sub>O (6 x 15 mL), brine  
 10 (20 mL) and dried (Na<sub>2</sub>SO<sub>4</sub>). Removal of solvent *in vacuo* then yielded crude product, which was purified by column chromatography using dichloromethane-methanol-aqueous ammonia (93:7:1) as eluent to furnish the product (110 mg, 57%).

<sup>1</sup>H-n.m.r. (CDCl<sub>3</sub>)  $\delta$  0.99 (t, 3H,  $J$ =7.4 Hz, CH<sub>3</sub>), 1.18 (t, 3H,  $J$ = 7.2 Hz, CH<sub>3</sub>), 1.33-1.60 (m, 2H, CH<sub>2</sub>), 1.82-2.02 (m, 2H, CH<sub>2</sub>), 2.11 (s, 3H, Ar-Me), 3.25-3.39 (m, 2H, CH<sub>2</sub>), 3.87 (s, 3H, OMe), 4.80-4.96 (m, 2H, 2 x NH), 5.26-5.36 (m, 1H, CH), 6.98 (br s, 1H, Ar-NHCONH),  
 15 7.22-7.28 (m, 1H, ArH), 7.67-7.72 (m, 2H, ArH), 7.83-7.88 (m, 1H, ArH), 8.05 (d, 1H,  $J$ = 0.8 Hz, ArH), 8.10 (d, 1H,  $J$ = 8.2 Hz, ArH), 8.47 (dd, 1H,  $J$ = 6.6, 1.8 Hz, ArH), 8.69 (d, 1H,  $J$ = 2.0 Hz, ArH).

### EXAMPLE 19

20 6-Chloro-N-(1-pyridin-3-ylbutyl)pyrimidin-4-amine



To a solution of 1-pyridin-3-ylbutan-1-amine (100 mg, 0.67 mmol) in ethanol (4 mL) was added 4,6-dichloropyrimidine (109 mg, 0.73 mmol) and diisopropylethylamine (232  $\mu$ L, 1.33 mmol). The solution was stirred for 3 days after which time the solvent was removed


49.

under reduced pressure. The residue was dissolved in ethyl acetate (25 mL) and washed with H<sub>2</sub>O (3 x 15 mL), brine (15 mL) and dried (Na<sub>2</sub>SO<sub>4</sub>). The residue remaining after concentration in vacuo was chromatographed using ethyl acetate-hexanes (4:6 – 9:1) as eluant to separate the desired product (28 mg).

- 5 <sup>1</sup>H-n.m.r. (CDCl<sub>3</sub>) δ 0.96 (t, 3H, *J* = 7.2 Hz, CH<sub>3</sub>), 1.22-1.51 (m, 2H, CH<sub>2</sub>), 1.76-1.96 (m, 2H, CH<sub>2</sub>), 4.76 (br s, 1H, CH), 5.63 (br s, 1H, NH), 6.23 (s, 1H, ArH), 7.26-7.32 (m, 1H, ArH), 7.59-7.65 (m, 1H, ArH), 8.33 (s, 1H, ArH), 8.53-8.56 (m, 1H, ArH), 8.60 (d, 1H, *J* = 1.8 Hz, ArH).

**EXAMPLE 20**

- 10 *2-Methoxy-4-{6-[(1-pyridin-3-ylbutyl)amino]pyrimidin-4-yl}phenol*



- Using an analogous procedure to that described in Example 12, reaction of 6-chloro-N-(1-pyridin-3-ylbutyl)pyrimidin-4-amine (24 mg, 0.09 mmol) with 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenol (27 mg, 0.11 mmol) afforded the desired product as a brown  
15 foam (12 mg).

- <sup>1</sup>H-n.m.r. (CDCl<sub>3</sub>) δ 0.97 (t, 3H, *J* = 7.2 Hz, CH<sub>3</sub>), 1.32-1.55 (m, 2H, CH<sub>2</sub>), 1.83-1.95 (m, 2H, CH<sub>2</sub>), 3.94 (s, 3H, OMe), 4.87 (br s, 1H, CH), 5.43 (br s, 1H, NH), 6.50 (s, 1H, ArH), 6.93 (d, 1H, *J* = 8.4 Hz, ArH), 7.25-7.31 (m, 2H, ArH), 7.55 (d, 1H, *J* = 2.0 Hz, ArH), 7.66-7.71 (m, 1H, ArH), 8.53 (dd, 1H, *J* = 4.9, 1.6 Hz, ArH), 8.58 (d, 1H, *J* = 1.6 Hz, ArH), 8.66 (d, 1H, *J* =  
20 2.0 Hz, ArH).

50.

**SCREENING****Compound Dilution**

For screening purposes, compounds were diluted in 96 well plates at a concentration of 20  $\mu$ M. Plates were warmed at 37°C for 30 minutes before assay.

**5 Growth and Maintenance of Cancer Cell lines**

K562 (Chronic Myeloid Leukemia), PC3 (Prostate Cancer), and DU145 (Prostate Cancer) were obtained from the American Type Culture Collection (ATCC). K562 was grown in RPMI, with 10% FBS with Glutamax added. DU145 cells were cultured in DMED, with 10% FBS and Glutamax and MEM non-essential amino acids added. PC3 cells were grown in F12K medium, with 10% FBS and Glutamax and MEM non-essential amino acids added. All cells were grown at 37°C in 5% CO<sub>2</sub>.

**Establishment of TEL:JAK cell lines**

The coding region encompassing nucleotides 1-487 of TEL was amplified by PCR using the oligonucleotides 5TEL (5' -GGA GGA TCC TGA TCT CTC TCG CTG TGA GAC-3') and 3TEL (5'-AGGC GTC GAC TTC TTC TTC ATG GTT CTG-3') and U937 mRNA as template. A BamH I site was present into the 5TEL Primer, a Sal I site was incorporated into the 3TEL primer. The regions encompassing the kinase domains of JAK2 (nucleotides 2994-3914; ; JAK2F 5'-ACGC GTC GAC GGT GCC TTT GAA GAC CGG GAT-3' ; JAK2R 5'-ATA GTT TAG CGG CCG CTC AGA ATG AAG GTC ATT T-3' ) and JAK3 (nucleotides 2520-3469; JAK3F 5'-GAA GTC GAC TAT GCC TGC CAA GAC CCC ACG ATC TT-3' ; JAK3R 5'-GGA TCT AGA CTA TGA AAA GGA CAG GGA GTG GTG TTT -3' ) were generated by PCR using Taq DNA Polymerase ( Gibco/BRL) and U937 mRNA as template. A SalI site was incorporated into the forward primer of JAK2 and JAK3, a Not I site was incorporated into the JAK2 reverse primer and a Xba I site was added to the reverse primer of JAK3.

A TEL/Jak2 fusion was generated by digestion of the TELPCR product with BamH I /Sal I, digestion of the JAK2 PCR product with Sal I/ Not I followed by ligation and subcloning into the mammalian expression Vector pTRE 2 (Clontech) digested with BamH I-Not I (pTELJAK2). For JAK3 Sal I/ Not I cleaved kinase domain PCR product was ligated with BamH I /Sal I cleaved TELproduct followed by ligation into BamH I/Not I cleaved pTRE2 (pTELJAK3).

51.

The growth factor dependent myelomonocytic cell line BaF3 bearing the pTET-off plasmid (Clontech) was transfected with either pTELJAK2 or pTELJAK3 and the cells selected for factor independent growth. BaF 3 wild type cells were cultured in DMEM 10% FCS, 10% WEHI 3B conditioned medium. BaF3 TELJAK cells were cultured in  
5 DMEM 10% Tet-System Approved FBS (without WEHI 3B conditioned medium).

**Cellular assays were performed as follows:**

Cell suspensions were prepared by harvesting cells from culture. (Cells used in this test should be in later log phase growth and high viability.) Cells were diluted in correct growth medium to 1.1x final concentration (from 50000 cell/mL to 200,000 cell/mL,  
10 depending on cell line).

Compounds to be tested were added (10µL, 10X final concentration) to a flat bottom 96-well plate. The cellular suspension (90µL per well) was added, and the plate incubated for 40 hr at 37 °C, 5% CO<sub>2</sub>. MTT (20 µL per well, 5mg/mL in PBS) was added and the plates were returned to the incubator for a further 6 hours. Lysis buffer (100 µL per well,  
15 10% SDS, 0.01N HCl) was added and the plate stored in the incubator overnight. The plate was then read at 590 nm.

**Tubulin Assay**

Turbidometric assays of microtubule assembly were performed by incubating microtubule protein in cuvettes at 37 °C in a thermostatically controlled  
20 spectrophotometer measuring the change in absorbance at 340nm over time. The microtubule protein was incubated with the each test compound at 0 °C and polymerisation was initiated by addition of 1mM GTP, prior to heating to 37 °C.







**Results**

The activity of a range of compounds is shown in Tables 1 and 2. Compounds that  
25 exhibited a capacity to inhibit greater than 50% of cellular growth at a concentration of 20µM are designated as "+"; compounds which did not inhibit 50% of cellular growth at 20µM are designated as "-"; compounds which were not tested are designated as "NT". Likewise, compounds which inhibited tubulin polymerisation by greater than 50% at 50µM are designated as "+"; compounds which did not inhibit tubulin polymerisation by  
30 50% at 50 µM are designated as "-"; and compounds which were not tested are designated as "NT".









52.







Table 1

| CHEMISTRY   | DU145 | PC3 | K562 | TelJak2 | TelJak3 | Tubulin |
|---|-------|-----|------|---------|---------|---------|
| <br>C24H28FN5O2  | +     | +   | +    | +       | +       | +       |
| <br>C27H37N5O2  | -     | -   | +    | +       | +       | NT      |
| <br>C21H23N3O  | +     | +   | +    | +       | +       | +       |
| <br>C26H32N4   | -     | -   | -    | +       | +       | NT      |
| <br>C24H29N3O3 | +     | +   | +    | +       | +       | +       |
| <br>C27H36N4O2 | +     | +   | +    | +       | +       | +       |







53.

| CHEMISTRY   | DU145 | PC3 | K562 | TelJak2 | TelJak3 | Tubulin |
|---|-------|-----|------|---------|---------|---------|
| <br>C24H29N5O2   | +     | +   | +    | +       | +       | +       |
| <br>C25H30N4O   | -     | -   | -    | +       | -       | NT      |
| <br>C26H33N5   | -     | -   | -    | +       | +       | NT      |
| <br>C23H28N6O2 | +     | +   | +    | +       | +       | +       |
| <br>C25H31N3O3 | +     | +   | +    | +       | +       | +       |
| <br>C28H36N4O3 | -     | +   | +    | +       | +       | NT      |







54.

| CHEMISTRY   | DU145 | PC3 | K562 | TelJak2 | TelJak3 | Tubulin |
|---|-------|-----|------|---------|---------|---------|
| <br>C21H24N4     | +     | +   | +    | +       | +       | +       |
| <br>C26H33N7O2  | -     | +   | -    | +       | +       | NT      |
| <br>C28H36N4O2 | +     | +   | +    | +       | +       | +       |
| <br>C28H38N4O2 | +     | +   | +    | +       | +       | +       |
| <br>C29H39N5O2 | +     | +   | +    | +       | +       | +       |
| <br>C27H31N5O2 | +     | +   | +    | +       | +       | +       |

55.







| CHEMISTRY   | DU145 | PC3 | K562 | TelJak2 | TelJak3 | Tubulin |
|---|-------|-----|------|---------|---------|---------|
| <br>C26H29N5O2   | -     | -   | +    | +       | +       | NT      |
| <br>C27H33N5O   | +     | +   | +    | +       | +       | +       |
| <br>C29H38N4O3 | -     | -   | +    | +       | +       | NT      |
| <br>C30H40N4O2 | +     | +   | +    | +       | +       | +       |
| <br>C29H38N4O2 | +     | +   | +    | +       | +       | +       |
| <br>C29H40N4O2 | +     | +   | +    | +       | +       | +       |

56.







| CHEMISTRY  | DU145 | PC3 | K562 | TelJak2 | TelJak3 | Tubulin |
|--|-------|-----|------|---------|---------|---------|
| <br>C30H41N5O2      | +     | +   | +    | +       | +       | +       |
| <br>C25H32N4O2     | +     | +   | +    | +       | +       | +       |
| <br>Chemistry 546 | -     | -   | +    | +       | +       | NT      |
| <br>C26H32N4O     | -     | -   | +    | +       | +       | NT      |
| <br>C26H32N4O     | -     | -   | +    | +       | +       | NT      |
| <br>C22H26N6O     | -     | -   | +    | +       | +       | NT      |









57.

| CHEMISTRY   | DU145 | PC3 | K562 | TelJak2 | TelJak3 | Tubulin |
|---|-------|-----|------|---------|---------|---------|
| <br>C26H26N6O2   | -     | -   | +    | +       | +       | NT      |
| <br>C27H33N5O   | +     | +   | +    | +       | +       | +       |
| <br>C21H24N4O2 | +     | +   | +    | +       | +       | +       |
| <br>C21H24N4O3 | -     | -   | -    | +       | +       | NT      |
| <br>C25H30N4   | +     | +   | +    | +       | +       | +       |
| <br>C21H25N5O  | +     | +   | +    | +       | +       | +       |







58.

| CHEMISTRY   | DU145 | PC3 | K562 | TelJak2 | TelJak3 | Tubulin |
|---|-------|-----|------|---------|---------|---------|
| <br>C22H26N6O2   | +     | +   | +    | +       | +       | +       |
| <br>C19H19N3O2  | +     | +   | +    | +       | +       | +       |
| <br>C24H28N2O3 | -     | +   | +    | +       | +       | NT      |
| <br>C20H20N2O2 | +     | +   | +    | +       | +       | +       |
| <br>C22H24N2O2 | +     | +   | +    | +       | +       | +       |
| <br>C23H26N2O  | +     | +   | +    | +       | +       | +       |







59.

| CHEMISTRY   | DU145 | PC3 | K562 | TelJak2 | TelJak3 | Tubulin |
|---|-------|-----|------|---------|---------|---------|
| <br>C26H32N6O2   | -     | -   | +    | +       | -       | NT      |
| <br>C25H31N5O   | -     | -   | -    | +       | +       | NT      |
| <br>C28H37N5O  | +     | -   | +    | +       | +       | NT      |
| <br>C28H35N5O2 | -     | -   | -    | +       | +       | NT      |
| <br>C26H31N5O  | NT    | -   | -    | +       | +       | NT      |
| <br>C25H28N4O2 | NT    | -   | -    | +       | +       | NT      |

60.







| CHEMISTRY   | DU145 | PC3 | K562 | TelJak2 | TelJak3 | Tubulin |
|---|-------|-----|------|---------|---------|---------|
| <br>C21H24N6O2   | NT    | -   | NT   | +       | +       | NT      |
| <br>C23H28N6O   | +     | +   | +    | +       | +       | +       |
| <br>C28H38N6O2 | -     | -   | -    | +       | +       | NT      |
| <br>C27H35N5O2 | -     | -   | +    | +       | +       | NT      |
| <br>C27H35N5O3 | -     | -   | -    | +       | +       | NT      |
| <br>C28H37N7O3 | +     | +   | +    | +       | +       | +       |

61.






| CHEMISTRY   | DU145 | PC3 | K562 | TelJak2 | TelJak3 | Tubulin |
|---|-------|-----|------|---------|---------|---------|
| <br>C25H33N7O2   | +     | +   | +    | +       | +       | +       |
| <br>C28H39N7O2   | -     | -   | -    | +       | +       | NT      |
| <br>C24H30N6O   | +     | +   | +    | +       | +       | +       |
| <br>C24H30N6O2 | +     | +   | +    | +       | +       | +       |
| <br>C23H28N6O2 | +     | +   | +    | +       | +       | +       |
| <br>C23H28N6O  | +     | +   | +    | +       | +       | +       |









62.

| CHEMISTRY   | DU145 | PC3 | K562 | TelJak2 | TelJak3 | Tubulin |
|---|-------|-----|------|---------|---------|---------|
| <br>C21H24N4O2   | -     | -   | +    | +       | +       | NT      |
| <br>C23H28N6O    | +     | +   | +    | +       | +       | +       |
| <br>C23H28N6O2  | +     | +   | +    | +       | +       | +       |
| <br>C24H30N6O2 | +     | +   | +    | +       | +       | +       |
| <br>C24H30N6O  | +     | +   | +    | +       | +       | +       |
| <br>C24H30N6O2 | +     | +   | +    | +       | +       | NT      |







63.

| CHEMISTRY   | DU145 | PC3 | K562 | TelJak2 | TelJak3 | Tubulin |
|---|-------|-----|------|---------|---------|---------|
| <br>C23H28N6O2   | +     | +   | +    | +       | +       | +       |
| <br>C24H30N6O2  | +     | +   | +    | +       | +       | +       |
| <br>C26H34N6O  | -     | -   | +    | +       | +       | NT      |
| <br>C22H26N6O2 | +     | +   | +    | +       | +       | +       |
| <br>C25H31N5O2 | +     | +   | +    | +       | +       | +       |







64.

| CHEMISTRY   | DU145 | PC3 | K562 | TelJak2 | TelJak3 | Tubulin |
|---|-------|-----|------|---------|---------|---------|
| <br>C24H29N5O3   | +     | +   | +    | +       | +       | +       |
| <br>C23H28N6O2  | +     | +   | +    | +       | +       | +       |
| <br>C21H23N3O3 | +     | -   | +    | +       | +       | NT      |
| <br>C23H27N5O3 | +     | +   | +    | +       | +       | +       |
| <br>C24H29N5O3 | +     | +   | +    | +       | +       | +       |
| <br>C20H22N4O2 | +     | +   | +    | +       | +       | +       |

65.



| CHEMISTRY   | DU145 | PC3 | K562 | TelJak2 | TelJak3 | Tubulin |
|---|-------|-----|------|---------|---------|---------|
| <br>C24H30N6O2   | +     | +   | +    | +       | +       | +       |
| <br>C21H24N4O2  | +     | +   | +    | +       | +       | +       |
| <br>C23H28N6O2 | +     | +   | +    | +       | +       | +       |
| <br>C20H22N4O2 | +     | +   | +    | +       | +       | +       |
| <br>C23H28N4O3 | -     | -   | +    | +       | +       | NT      |
| <br>C19H20N4O2 | -     | -   | +    | +       | +       | NT      |

66.

| CHEMISTRY   | DU145 | PC3 | K562 | TelJak2 | TelJak3 | Tubulin |
|---|-------|-----|------|---------|---------|---------|
| <br>C24H30N6O2   | +     | +   | +    | +       | +       | +       |
| <br>C21H24N4O2  | +     | +   | +    | +       | +       | +       |
| <br>C23H28N6O2 | +     | +   | +    | +       | +       | +       |
| <br>C20H22N4O2 | +     | +   | +    | +       | +       | +       |
| <br>C23H28N4O3 | -     | -   | +    | +       | +       | NT      |
| <br>C19H20N4O2 | -     | -   | +    | +       | +       | NT      |









67.

| CHEMISTRY  | DU145 | PC3 | K562 | TelJak2 | TelJak3 | Tubulin |
|--|-------|-----|------|---------|---------|---------|
| <br>C23H28N6O2  | +     | +   | +    | +       | +       | +       |
| <br>C19H19N3O2 | +     | +   | +    | +       | +       | +       |

68.

Table 2

| Chemistry   | TelJak2 | TelJak3 | DU145 | Tubulin |
|---|---------|---------|-------|---------|
| <br>C25H23N5O3   | +       | +       | +     | +       |
| <br>C24H22N4O2 | +       | +       | +     | +       |
| <br>C24H27N5O3 | +       | +       | +     | +       |
| <br>C25H23N5O3 | +       | +       | +     | +       |
| <br>C20H21N3O2 | +       | +       | +     | +       |
| <br>C19H19N3O2 | +       | +       | +     | +       |

69.

Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

- 5 All publications mentioned in this specification are herein incorporated by reference. Any discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is solely for the purpose of providing a context for the present invention. It is not to be taken as an admission that any or all of these matters form part of the prior art base or were common general knowledge in the field relevant to the present invention as it existed anywhere before the priority date of each claim of this application.
- 10

It will be appreciated by persons skilled in the art that numerous variations and/or modifications may be made to the invention as shown in the specific embodiments without departing from the spirit or scope of the invention as broadly described. The present embodiments are, therefore, to be considered in all respects as illustrative and not restrictive.

70.


## REFERENCES

1. Jordan M A. and Wilson L. (1998) Microtubules and actin filaments: dynamic targets for cancer chemotherapy. *Curr. Op. Cell Biol.*, 10, 123-130.
2. Jordan MA, Margolis RL, Himes RH, Wilson L., (1986) Identification of a distinct class  
5 of vinblastine binding sites on microtubules. *J. Mol. Biol.* 187:61-73
3. Rai SS, and Wolff J., (1996) Localization of the vinblastine-binding site on  $\beta$ -tubulin. *J. Biol. Chem.* 271:14707-11

71.

## CLAIMS

1. A method of modulating microtubule polymerisation in a subject, said method comprising administering a therapeutically effective amount of at least one compound of the general formula (I)



I

or pharmaceutically acceptable prodrugs, salts, hydrates, solvates, crystal forms or diastereomers thereof, wherein:

R1 is H, C<sub>1-4</sub> alkyl;

Q is a bond, or C<sub>1-4</sub> alkyl;

A is aryl, hetaryl optionally substituted with 0-3 substituents independently chosen from halogen, C<sub>1-4</sub> alkyl, CH<sub>2</sub>F, CHF<sub>2</sub>, CF<sub>3</sub>, CN, aryl, hetaryl, OCF<sub>3</sub>, OC<sub>1-4</sub>alkyl, OC<sub>2-5</sub>alkylNR<sub>4</sub>R<sub>5</sub>, Oaryl, Ohetaryl, CO<sub>2</sub>R<sub>4</sub>, CONR<sub>4</sub>R<sub>5</sub>, nitro, NR<sub>4</sub>R<sub>5</sub>, C<sub>1-4</sub> alkylNR<sub>4</sub>R<sub>5</sub>, NR<sub>6</sub>C<sub>1-4</sub>alkylNR<sub>4</sub>R<sub>5</sub>, NR<sub>4</sub>COR<sub>5</sub>, NR<sub>6</sub>CONR<sub>4</sub>R<sub>5</sub>, NR<sub>4</sub>SO<sub>2</sub>R<sub>5</sub>;

R<sub>4</sub>, R<sub>5</sub> are each independently H, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl cycloalkyl, C<sub>1-4</sub> alkyl cyclohetalkyl, aryl, hetaryl, C<sub>1-4</sub>alkyl aryl, C<sub>1-4</sub> alkyl hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>7</sub>;

R<sub>6</sub> is selected from H, C<sub>1-4</sub> alkyl;

R<sub>7</sub> is selected from H, C<sub>1-4</sub> alkyl, aryl, hetaryl, C<sub>1-4</sub> alkyl aryl, C<sub>1-4</sub> alkyl hetaryl;

R<sub>2</sub> is 0-2 substituents independently selected from halogen, C<sub>1-4</sub>alkyl, OH, OC<sub>1-4</sub>alkyl, CH<sub>2</sub>F, CHF<sub>2</sub>, CF<sub>3</sub>, OCF<sub>3</sub>, CN, C<sub>1-4</sub>alkylNR<sub>8</sub>R<sub>9</sub>, OC<sub>1-4</sub>alkylNR<sub>8</sub>R<sub>9</sub>, CO<sub>2</sub>R<sub>8</sub>, CONR<sub>8</sub>R<sub>9</sub>, NR<sub>8</sub>R<sub>9</sub>, NR<sub>8</sub>COR<sub>9</sub>, NR<sub>10</sub>CONR<sub>8</sub>R<sub>9</sub>, NR<sub>8</sub>SO<sub>2</sub>R<sub>9</sub>;



72.

R8, R9 are each independently H, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl cycloalkyl, C<sub>1-4</sub> alkyl cyclohetalkyl, aryl, hetaryl, C<sub>1-4</sub> alkyl aryl, C<sub>1-4</sub> alkyl hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR11;

R10 is selected from H, C<sub>1-4</sub> alkyl, aryl or hetaryl;

R11 is selected from H, C<sub>1-4</sub> alkyl, aryl, hetaryl, C<sub>1-4</sub> alkyl aryl, C<sub>1-4</sub> alkyl hetaryl;

Y is halogen, OH, NR12R13, NR14COR12, NR14CONR12R13, N14SO<sub>2</sub>R13;

R12 and R13 are each independently H, CH<sub>2</sub>F, CHF<sub>2</sub>, CF<sub>3</sub>, CN, C<sub>1-4</sub> alkyl optionally substituted with OH, OC<sub>1-4</sub>alkyl or NR15R16, cycloalkyl; cyclohetalkyl, C<sub>1-4</sub> alkyl cycloalkyl, C<sub>1-4</sub> alkyl cyclohetalkyl, or may be joined to form an optionally substituted 3-6 membered ring optionally containing an atom selected from O, S, NR14

R14, R15 and R16 are each independently selected from H, C<sub>1-4</sub> alkyl;


n = 0-4;

W is selected from H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl; where C<sub>1-4</sub>alkyl or C<sub>2-6</sub>alkenyl may be optionally substituted with C<sub>1-4</sub>alkyl, OH, OC<sub>1-4</sub>alkyl, NR15R16;


R15, and R16 are each independently H, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl cycloalkyl, C<sub>1-4</sub> alkyl cyclohetalkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR17;

R17 is selected from H, C<sub>1-4</sub> alkyl.


2. A method according to claim 1 wherein the compound is selected from the group consisting of:



73.



3. A method according to claim 1 or claim 2, wherein said method is used in the treatment of a hyperproliferation-related disorder or disease state.
4. A method according to claim 2, wherein the hyperproliferation-related disorder or disease state is selected from the group consisting of Cancer, infectious diseases, vascular restenosis and inflammatory diseases.
5. A compound of the general formula (II)



II

or pharmaceutically acceptable prodrugs, salts, hydrates, solvates, crystal forms or diastereomers thereof, wherein:

74.

R1 is H, C<sub>1-4</sub> alkyl;

Q is a bond, or C<sub>1-4</sub> alkyl;

A is aryl, hetaryl optionally substituted with 0-3 substituents independently chosen from halogen, C<sub>1-4</sub> alkyl, CH<sub>2</sub>F, CHF<sub>2</sub>, CF<sub>3</sub>, CN, aryl, hetaryl, OCF<sub>3</sub>, OC<sub>1-4</sub>alkyl, OC<sub>2-5</sub>alkylNR<sub>4</sub>R<sub>5</sub>, Oaryl, Ohetaryl, CO<sub>2</sub>R<sub>4</sub>, CONR<sub>4</sub>R<sub>5</sub>, nitro, NR<sub>4</sub>R<sub>5</sub>, C<sub>1-4</sub> alkylNR<sub>4</sub>R<sub>5</sub>, NR<sub>6</sub>C<sub>1-4</sub>alkylNR<sub>4</sub>R<sub>5</sub>, NR<sub>4</sub>COR<sub>5</sub>, NR<sub>6</sub>CONR<sub>4</sub>R<sub>5</sub>, NR<sub>4</sub>SO<sub>2</sub>R<sub>5</sub>;

R<sub>4</sub>, R<sub>5</sub> are each independently H, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl cycloalkyl, C<sub>1-4</sub> alkyl cyclohetalkyl, aryl, hetaryl, C<sub>1-4</sub>alkyl aryl, C<sub>1-4</sub> alkyl hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>7</sub>;







R<sub>6</sub> is selected from H, C<sub>1-4</sub> alkyl;

R<sub>7</sub> is selected from H, C<sub>1-4</sub> alkyl, aryl, hetaryl, C<sub>1-4</sub> alkyl aryl, C<sub>1-4</sub> alkyl hetaryl;


R<sub>2</sub> is 0-2 substituents independently selected from C<sub>1-4</sub>alkyl and OC<sub>1-4</sub>alkyl;

Y is CH<sub>2</sub>OH, OC<sub>1-4</sub>alkylOH, OC<sub>1-4</sub>alkylR<sub>12</sub>, OC<sub>1-4</sub>alkylNR<sub>12</sub>NR<sub>13</sub>, C(O)R<sub>12</sub>, CH<sub>2</sub>R<sub>12</sub>, COOR<sub>12</sub>, CONR<sub>12</sub>R<sub>13</sub>, OCONR<sub>12</sub>R<sub>13</sub>, CH<sub>2</sub>NR<sub>12</sub>R<sub>13</sub>, NHCOR<sub>12</sub>, NHCONR<sub>12</sub>R<sub>13</sub>,

R<sub>12</sub> and R<sub>13</sub> are each independently H, C<sub>1-2</sub> alkyl, (CH<sub>2</sub>)<sub>3</sub>NEt<sub>2</sub>, (CH<sub>2</sub>)<sub>2</sub>NMe<sub>2</sub>, (CH<sub>2</sub>)<sub>5</sub>NH<sub>2</sub>, (CH<sub>2</sub>)<sub>2</sub>OH,



75.



n = 0-4;


W is selected from H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl; where C<sub>1-4</sub>alkyl or C<sub>2-6</sub>alkenyl may be optionally substituted with C<sub>1-4</sub>alkyl, OH, OC<sub>1-4</sub>alkyl, NR<sub>15</sub>R<sub>16</sub>;

R<sub>15</sub>, and R<sub>16</sub> are each independently H, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl cycloalkyl, C<sub>1-4</sub> alkyl cyclohetalkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>17</sub>


R<sub>17</sub> is selected from H, C<sub>1-4</sub> alkyl;

wherein when Y is CH<sub>2</sub>R<sub>12</sub> then R<sub>12</sub> is not H, C<sub>1-2</sub>alkyl.













6. A compound according to claim 5 selected from the group consisting of:




76.



77.




78.







79.




80.



81.



7. A compound of the general formula (III)



III

or pharmaceutically acceptable prodrugs, salts, hydrates, solvates, crystal forms or diastereomers thereof, wherein:

$X_1, X_2, X_3, X_4$  are selected from the following:

- (i)  $X_1$  and  $X_2$  are N and  $X_3$  and  $X_4$  are C independently substituted with Y;
- (ii)  $X_1$  and  $X_4$  are N and  $X_2$  and  $X_3$  are C independently substituted with Y;
- (iii)  $X_1$  and  $X_3$  are N and  $X_2$  and  $X_4$  are C independently substituted with Y;
- (iv)  $X_2$  and  $X_4$  are N and  $X_1$  and  $X_3$  are C independently substituted with Y;
- (v)  $X_1$  is N and  $X_2, X_3$ , and  $X_4$  are C independently substituted with Y;

82.

- (vi)  $X_3$  is N and  $X_1$ ,  $X_2$ , and  $X_4$  are C independently substituted with Y;
- (vii)  $X_4$  is N and  $X_1$ ,  $X_2$ , and  $X_3$  are C independently substituted with Y;
- (viii)  $X_2$  is N and  $X_1$ ,  $X_3$ , and  $X_4$  are C independently substituted with Y; and
- (ix)  $X_1$ ,  $X_2$  and  $X_3$  are N and  $X_4$  is C substituted with Y;

$R_1$  is H,  $C_{1-6}$ alkyl,  $C_{1-6}$  alkylNR<sub>5</sub>R<sub>6</sub>,  $C_{1-6}$ alkylNR<sub>5</sub>COR<sub>6</sub>,  $C_{1-6}$ alkylNR<sub>5</sub>SO<sub>2</sub>R<sub>6</sub>,  $C_{1-6}$ alkylCO<sub>2</sub>R<sub>5</sub>,  $C_{1-6}$ alkylCONR<sub>5</sub>R<sub>6</sub>, where R<sub>5</sub> and R<sub>6</sub> are each independently H,  $C_{1-4}$ alkyl, aryl, hetaryl,  $C_{1-4}$ alkylaryl,  $C_{1-4}$ alkylhetaryl or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>7</sub>;

$R_7$  is selected from H,  $C_{1-4}$ alkyl;

$R_2$  is selected from  $C_{1-6}$ alkylOH, OC<sub>2-6</sub>alkylOH,  $C_{1-6}$ alkylNR<sub>8</sub>R<sub>9</sub>, OC<sub>2-6</sub>alkylNR<sub>8</sub>R<sub>9</sub>,  $C_{1-6}$ alkylNR<sub>8</sub>COR<sub>9</sub>, OC<sub>2-6</sub>alkylNR<sub>8</sub>COR<sub>9</sub>,  $C_{1-6}$ alkylhetaryl, OC<sub>2-6</sub>alkylhetaryl, OCONR<sub>8</sub>R<sub>9</sub>, NR<sub>8</sub>COOR<sub>9</sub>, NR<sub>10</sub>CONR<sub>8</sub>R<sub>9</sub>, CONR<sub>8</sub>R<sub>9</sub>, NR<sub>8</sub>COR<sub>12</sub>;

$R_8$ ,  $R_9$  are each independently H,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkylNR<sub>11</sub>R<sub>13</sub>, hetaryl, cyclohetalkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>14</sub>;

$R_{12}$  is  $C_{2-4}$ alkyl,  $C_{1-4}$ alkylNR<sub>11</sub>R<sub>13</sub>, hetaryl, cyclohetalkyl;

$R_{11}$ ,  $R_{13}$  are each independently H,  $C_{1-4}$ alkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>14</sub>;

$R_{14}$  is selected from H,  $C_{1-4}$ alkyl;

$R_{10}$  is H,  $C_{1-4}$  alkyl;

$R_3$  and  $R_4$  are each independently H, halogen,  $C_{1-4}$ alkyl, OH, OC<sub>1-4</sub>alkyl, CF<sub>3</sub>, OCF<sub>3</sub>;

$Q$  is a bond, or  $C_{1-4}$  alkyl;

$W$  is selected from H,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl; where  $C_{1-4}$ alkyl or  $C_{2-6}$ alkenyl may be optionally substituted with  $C_{1-4}$ alkyl, OH, OC<sub>1-4</sub>alkyl, NR<sub>15</sub>R<sub>16</sub>;

83.

R15, and R16 are each independently H, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyl cycloalkyl, C<sub>1-4</sub>alkyl cyclohetalkyl, aryl, hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR17;

R17 is selected from H, C<sub>1-4</sub>alkyl;

A is aryl, hetaryl optionally substituted with 0-3 substituents independently chosen from halogen, C<sub>1-4</sub> alkyl, CF<sub>3</sub>, aryl, hetaryl, OCF<sub>3</sub>, OC<sub>1-4</sub>alkyl, OC<sub>2-5</sub>alkylNR18R19, Oaryl, Ohetaryl, CO<sub>2</sub>R18, CONR18R19, NR18R19, C<sub>1-4</sub> alkylNR18R19, NR20C<sub>1-4</sub>alkylNR18R19, NR18COR19, NR20CONR18R19, NR18SO<sub>2</sub>R19;

R18, R19 are each independently H, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl cyclohetalkyl, aryl, hetaryl, C<sub>1-4</sub>alkyl aryl, C<sub>1-4</sub> alkyl hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR21;


R21 is selected from H, C<sub>1-4</sub>alkyl;

R20 is selected from H, C<sub>1-4</sub>alkyl;

Y is selected from H, C<sub>1-4</sub>alkyl, OH, NR22R23;

R22, R23 are each independently H, C<sub>1-4</sub>alkyl.

8. A compound according to formula (III) of claim 7, wherein the compound is of the general formula (IV)



IV

or pharmaceutically acceptable prodrugs, salts, hydrates, solvates, crystal forms or diastereomers thereof, wherein:

X<sub>1</sub>, X<sub>2</sub>, X<sub>3</sub>, X<sub>4</sub> are selected from the following:

84.

- (i)  $X_1$  and  $X_2$  are N and  $X_3$  and  $X_4$  are C independently substituted with Y;
- (ii)  $X_1$  and  $X_4$  are N and  $X_2$  and  $X_3$  are C independently substituted with Y;
- (iii)  $X_1$  and  $X_3$  are N and  $X_2$  and  $X_4$  are C independently substituted with Y;
- (iv)  $X_2$  and  $X_4$  are N and  $X_1$  and  $X_3$  are C independently substituted with Y;
- (v)  $X_1$  is N and  $X_2$ ,  $X_3$ , and  $X_4$  are C independently substituted with Y;
- (vi)  $X_3$  is N and  $X_1$ ,  $X_2$ , and  $X_4$  are C independently substituted with Y;
- (vii)  $X_4$  is N and  $X_1$ ,  $X_2$ , and  $X_3$  are C independently substituted with Y;
- (viii)  $X_2$  is N and  $X_1$ ,  $X_3$ , and  $X_4$  are C independently substituted with Y; and
- (ix)  $X_1$ ,  $X_2$  and  $X_3$  are N and  $X_4$  is C substituted with Y;

$R_1$  is H,  $C_{1-6}$ alkyl,  $C_{1-6}$  alkylNR<sub>5</sub>R<sub>6</sub>, where  $R_5$  and  $R_6$  are each independently H,  $C_{1-4}$ alkyl, aryl, hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>7</sub>;

$R_7$  is selected from H,  $C_{1-4}$  alkyl;

$R_2$  is selected from  $C_{1-6}$ alkylOH,  $OC_{2-6}$ alkylOH,  $C_{1-6}$ alkylNR<sub>8</sub>R<sub>9</sub>,  $OC_{2-6}$ alkylNR<sub>8</sub>R<sub>9</sub>,  $C_{1-6}$ alkylNR<sub>8</sub>COR<sub>9</sub>,  $OC_{2-6}$ alkylNR<sub>8</sub>COR<sub>9</sub>,  $C_{1-6}$ alkylhetaryl,  $OC_{2-6}$ alkylhetaryl,  $OCONR_8R_9$ ,  $NR_8COOR_9$ ,  $NR_{10}CONR_8R_9$ ,  $CONR_8R_9$ ,  $NR_8COR_{12}$ ;

$R_8$ ,  $R_9$  are each independently H,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkylNR<sub>11</sub>R<sub>13</sub>, hetaryl, cyclohetalkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>14</sub>;

$R_{12}$  is  $C_{2-4}$ alkyl,  $C_{1-4}$ alkylNR<sub>11</sub>R<sub>13</sub>, hetaryl, cyclohetalkyl;

$R_{11}$ ,  $R_{13}$  are each independently H,  $C_{1-4}$ alkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>14</sub>;

$R_{14}$  is selected from H,  $C_{1-4}$ alkyl;

$R_{10}$  is H,  $C_{1-4}$ alkyl;

85.

R3 and R4 are each independently H, halogen, C<sub>1-4</sub>alkyl, OH, OC<sub>1-4</sub>alkyl, CF<sub>3</sub>, OCF<sub>3</sub>;

Q is CH;

W is selected from C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl; where C<sub>1-4</sub>alkyl or C<sub>2-6</sub>alkenyl may be optionally substituted with C<sub>1-4</sub>alkyl, OH, OC<sub>1-4</sub>alkyl, NR<sub>15</sub>R<sub>16</sub>;

R<sub>15</sub>, and R<sub>16</sub> are each independently H, C<sub>1-4</sub>alkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>17</sub>;

R<sub>17</sub> is selected from H, C<sub>1-4</sub>alkyl;

A is aryl, hetaryl optionally substituted with 0-2 substituents independently chosen from halogen, C<sub>1-4</sub>alkyl, CF<sub>3</sub>, aryl, hetaryl, OCF<sub>3</sub>, OC<sub>1-4</sub>alkyl, OC<sub>2-5</sub>alkyl, NR<sub>18</sub>R<sub>19</sub>, Oaryl, Ohetaryl, CO<sub>2</sub>R<sub>18</sub>, CONR<sub>18</sub>R<sub>19</sub>, NR<sub>18</sub>R<sub>19</sub>, C<sub>1-4</sub>alkyl, NR<sub>18</sub>R<sub>19</sub>, NR<sub>20</sub>C<sub>1-4</sub>alkyl, NR<sub>18</sub>R<sub>19</sub>, NR<sub>18</sub>COR<sub>19</sub>, NR<sub>20</sub>CONR<sub>18</sub>R<sub>19</sub>, NR<sub>18</sub>SO<sub>2</sub>R<sub>19</sub>;

R<sub>18</sub>, R<sub>19</sub> are each independently H, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyl cyclohetalkyl, aryl, hetaryl, C<sub>1-4</sub>alkyl aryl, C<sub>1-4</sub>alkyl hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>21</sub>;


R<sub>21</sub> is selected from H, C<sub>1-4</sub>alkyl;

R<sub>20</sub> is selected from H, C<sub>1-4</sub>alkyl;

Y is selected from H, C<sub>1-4</sub>alkyl, NR<sub>22</sub>R<sub>23</sub>;


R<sub>22</sub>, R<sub>23</sub> are each independently H, C<sub>1-4</sub>alkyl.

9. A compound according to claim 7 wherein the compound is selected from the group consisting of:






86.



10. A compound of the general formula (V)



or pharmaceutically acceptable prodrugs, salts, hydrates, solvates, crystal forms or diastereomers thereof, wherein:

$X_1, X_2, X_3, X_4$  are selected from the following:

- (i)  $X_1$  and  $X_2$  are N and  $X_3$  and  $X_4$  are C independently substituted with Y;

87.

- (ii)  $X_1$  and  $X_4$  are N and  $X_2$  and  $X_3$  are C independently substituted with Y;
- (iii)  $X_2$  and  $X_4$  are N and  $X_1$  and  $X_3$  are C independently substituted with Y;
- (iv)  $X_1$  is N and  $X_2$ ,  $X_3$ , and  $X_4$  are C independently substituted with Y;
- (v)  $X_3$  is N and  $X_1$ ,  $X_2$ , and  $X_4$  are C independently substituted with Y;
- (vi)  $X_4$  is N and  $X_1$ ,  $X_2$ , and  $X_3$  are C independently substituted with Y;
- (vii)  $X_2$  is N and  $X_1$ ,  $X_3$ , and  $X_4$  are C independently substituted with Y; and
- (viii)  $X_1$ ,  $X_2$  and  $X_3$  are N and  $X_4$  is C substituted with Y;

R1 is H,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkylNR5R6,  $C_{1-6}$ alkylNR5COR6,  $C_{1-6}$ alkylNR5SO<sub>2</sub>R6,  $C_{1-6}$ alkylCO<sub>2</sub>R5,  $C_{1-6}$ alkylCONR5R6, where R5 and R6 are each independently H,  $C_{1-4}$ alkyl, aryl, hetaryl,  $C_{1-4}$ alkylaryl,  $C_{1-4}$ alkylhetaryl or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR7;

R7 is selected from H,  $C_{1-4}$ alkyl;

R2 is selected from OH, OC<sub>1-6</sub>alkyl,  $C_{1-6}$ alkylOH, OC<sub>2-6</sub>alkylOH,  $C_{1-6}$ alkylNR8R9, OC<sub>2-6</sub>alkylNR8R9,  $C_{1-6}$ alkylNR8COR9, OC<sub>2-6</sub>alkylNR8COR9,  $C_{1-6}$ alkylhetaryl, OC<sub>2-6</sub>alkylhetaryl, OCONR8R9, NR8COOR9, NR10CONR8R9, CONR8R9, NR8COR12;

R8, R9 are each independently H,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkylNR11R13, hetaryl, cyclohetalkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR14;

R12 is  $C_{2-4}$ alkyl,  $C_{1-4}$ alkylNR11R13, hetaryl, cyclohetalkyl;

R11, R13 are each independently H,  $C_{1-4}$ alkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR14;

R14 is selected from H,  $C_{1-4}$ alkyl;

R10 is H,  $C_{1-4}$ alkyl;

R3 and R4 are each independently H, halogen,  $C_{1-4}$ alkyl, OH, OC<sub>1-4</sub>alkyl, CF<sub>3</sub>, OCF<sub>3</sub>;

88.

Q is a bond, or C<sub>1-4</sub>alkyl;

W is selected from H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl; where C<sub>1-4</sub>alkyl or C<sub>2-6</sub>alkenyl may be optionally substituted with C<sub>1-4</sub>alkyl, OH, OC<sub>1-4</sub>alkyl, NR<sub>15</sub>R<sub>16</sub>;

R<sub>15</sub>, and R<sub>16</sub> are each independently H, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyl cycloalkyl, C<sub>1-4</sub>alkyl cyclohetalkyl, aryl, hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>17</sub>;

R<sub>17</sub> is selected from H, C<sub>1-4</sub>alkyl;

A is aryl, hetaryl optionally substituted with 0-3 substituents independently chosen from halogen, C<sub>1-4</sub> alkyl, CF<sub>3</sub>, aryl, hetaryl, OCF<sub>3</sub>, OC<sub>1-4</sub>alkyl, OC<sub>2-5</sub>alkylNR<sub>18</sub>R<sub>19</sub>, Oaryl, Ohetaryl, CO<sub>2</sub>R<sub>18</sub>, CONR<sub>18</sub>R<sub>19</sub>, NR<sub>18</sub>R<sub>19</sub>, C<sub>1-4</sub> alkylNR<sub>18</sub>R<sub>19</sub>, NR<sub>20</sub>C<sub>1-4</sub>alkylNR<sub>18</sub>R<sub>19</sub>, NR<sub>18</sub>COR<sub>19</sub>, NR<sub>20</sub>CONR<sub>18</sub>R<sub>19</sub>, NR<sub>18</sub>SO<sub>2</sub>R<sub>19</sub>;

R<sub>18</sub>, R<sub>19</sub> are each independently H, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl cyclohetalkyl, aryl, hetaryl, C<sub>1-4</sub>alkyl aryl, C<sub>1-4</sub> alkyl hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR<sub>21</sub>;


R<sub>21</sub> is selected from H, C<sub>1-4</sub> alkyl;

R<sub>20</sub> is selected from H, C<sub>1-4</sub> alkyl;


Y is selected from H, C<sub>1-4</sub>alkyl, OH, NR<sub>22</sub>R<sub>23</sub>;

R<sub>22</sub>, R<sub>23</sub> are each independently H, C<sub>1-4</sub> alkyl.

11. A compound according to claim 10 selected from the group consisting of:




89.



90.


12. A compound of the formula:



or a pharmaceutically acceptable prodrug, salt, hydrate, solvate, crystal form or a diastereomer thereof

91.

13. A compound of the formula:



or a pharmaceutically acceptable prodrug, salt, hydrate, solvate, crystal form or a diastereomer thereof

14. A composition comprising a carrier and at least one compound according to any one of claims 5 to 13.

15. A method of treatment of a hyperproliferation-related disorder or disease state in a subject, said method comprising administering a therapeutically effective amount of at least one compound according to any one claims 1 to 13 or a composition according to 14.

16. A method of treatment according to claim 15, wherein the hyperproliferation-related disorder or disease state is treatable by the modulation of microtubule polymerisation.

17. A method according to claim 15 or claim 16, wherein the hyperproliferation-related disorder or disease state is selected from the group consisting of Cancer, infectious diseases, vascular restenosis or inflammatory diseases.

18. A method of treatment of a protein-kinase related disorder or disease state in a subject, said method comprising administering a therapeutically effective amount of at least one compound according to any one of claims 1 to 13 or a composition according to 14.

17. A method according to claim 18, wherein the protein-kinase related disorder or disease state is selected from the group consisting of Atopy, Cell Mediated Hypersensitivity, Rheumatic Diseases, Other autoimmune diseases and Viral Diseases.

18. A method of treatment of diseases and conditions associated with inflammation and infection in a subject, said method comprising administering a therapeutically effective amount of at least one compound according to any one of claims 1 to 13 or a composition according to claim 14.

# INTERNATIONAL SEARCH REPORT

International application No.

**PCT/AU2004/001689**

## A. CLASSIFICATION OF SUBJECT MATTER

Int. Cl. <sup>7</sup>: C07D 213/74, 237/20, 239/42, 241/20, 253/07, 401/12, 401/14, 413/10, 403/10; A61K 31/4418, 31/444, 31/496, 31/497, 31/506, 31/53, 31/5377; A61P 29/00, 37/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

**SEE ELECTRONIC DATABASES BELOW**

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN Files Registry, CA: substructure search based on claim 7, limited by examples

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages  | Relevant to claim No.    |
|-----------|---|--------------------------|
| P,X       | WO 2004/052868 (CYTOPIA PTY. LTD.) 24 June 2004<br>See whole document, Formulae I and II, Table 1, p. 11 (RH column, 3 <sup>rd</sup> compound), p. 59 (C23H25FN4O2, C28H37FN4O), p. 63 (C23H27N5O2, C22H25N5O2), p. 64 (C24H28N5O2, C23H26N4O2) | 1-9, 12, 14-18           |
| P,X       | WO 2003/099796 (CYTOPIA PTY. LTD.) 4 December 2003<br>See Formulae I and II, Example 35   | 5-10, 12, 14, 15, 17, 18 |
| X         | WO 2003/031406 (IRM LLC) 17 April 2003<br>See Tables 2-4, Entries 1-5 and 12  | 5, 10, 14, 18            |



Further documents are listed in the continuation of Box C



See patent family annex

|          |   |     |  |
|----------|---|-----|--|
| *<br>"A" | Special categories of cited documents:<br>document defining the general state of the art which is not considered to be of particular relevance                      | "T" | later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention  |
| "E"      | earlier application or patent but published on or after the international filing date   | "X" | document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone   |
| "L"      | document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) | "Y" | document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art |
| "O"      | document referring to an oral disclosure, use, exhibition or other means  | "&" | document member of the same patent family  |
| "P"      | document published prior to the international filing date but later than the priority date claimed  |     |  |

Date of the actual completion of the international search  
**14 January 2005**

Date of mailing of the international search report

**3 FEB 2005**

Name and mailing address of the ISA/AU

AUSTRALIAN PATENT OFFICE  
PO BOX 200, WODEN ACT 2606, AUSTRALIA  
E-mail address: [pct@ipaaustralia.gov.au](mailto:pct@ipaaustralia.gov.au)  
Facsimile No. (02) 6285 3929

Authorized officer

**L.F. MCCAFFERY**

Telephone No : (02) 6283 2573



# INTERNATIONAL SEARCH REPORT

International application No.  
**PCT/AU2004/001689**

| C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT |  |                            |
|---|--|----------------------------|
| Category*   | Citation of document, with indication, where appropriate, of the relevant passages   | Relevant to claim No.      |
| X   | WO 2003/026661 (YAMANOUCHI PHARMACEUTICAL CO. LTD) 3 April 2003<br>See Abstract, Examples 3, 6, 7 and 346                                | 7, 8, 10                   |
| X   | WO 2002/060492 (CYTOPIA PTY. LTD.) 8 August 2002<br>See whole document, Example 4, compounds 20310, 20504, 20506, 20508, 25210 and 32102 | 5-9, 12, 14,<br>15, 17, 18 |

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

**PCT/AU2004/001689**

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

| Patent Document Cited in<br>Search Report   |          | Patent Family Member |         |    |            |
|---|----------|----------------------|---------|----|------------|
| WO  | 04052868 |                      |         |    |            |
| WO  | 03099796 | GB                   | 2392154 | US | 2004235862 |
| WO  | 03031406 | CA                   | 2463563 | US | 2003191312 |
| WO  | 03026661 |                      |         |    |            |
| WO  | 02060492 | CA                   | 2436487 | EP | 1363702    |
|   |          |                      |         | US | 2004102455 |
| Due to data integration issues this family listing may not include 10 digit Australian applications filed since May 2001. |          |                      |         |    |            |
| END OF ANNEX  |          |                      |         |    |            |